


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 239/00	A2	(11) International Publication Number: WO 98/45276 (43) International Publication Date: 15 October 1998 (15.10.98)
(21) International Application Number: PCT/US98/06733 (22) International Filing Date: 7 April 1998 (07.04.98) (30) Priority Data: 08/838,540 9 April 1997 (09.04.97) US 60/071,322 14 January 1998 (14.01.98) US (71) Applicant: DU PONT PHARMACEUTICALS COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors: CORBETT, Jeffrey, W.; 1639 Oak Avenue, Haddon Height, NJ 08035 (US). KO, Soo, Sung; 7 Aston Circle, Hockessin, DE 19707 (US). (74) Agent: VANCE, David, H.; Du Pont Pharmaceuticals Com- pany, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		(81) Designated States: AM, AU, AZ, BR, BY, CA, CN, CZ, EE, HU, IL, JP, KG, KR, KZ, LT, LV, MD, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, UA, VN, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: 4,4-DISUBSTITUTED-3,4-DIHYDRO-2(1H)-QUINAZOLINONES USEFUL AS HIV REVERSE TRANSCRIPTASE IN- HIBITORS <div data-bbox="743 1182 1159 1297"></div> (57) Abstract <p>The present invention relates to 4,4-disubstituted-3,4-dihydro-2(1H)-quinazolines of formula (I), or stereoisomeric forms, stereoisomeric mixtures, or pharmaceutically acceptable salt forms thereof, which are useful as inhibitors of HIV reverse transcriptase, and to pharmaceutical compositions and diagnostic kits comprising the same, and methods of using the same for treating viral infection or as an assay standard or reagent.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE

4,4-DISUBSTITUTED-3,4-DIHYDRO-2(1H)-QUINAZOLINONES USEFUL AS
HIV REVERSE TRANSCRIPTASE INHIBITORS

5

FIELD OF THE INVENTION

This invention relates generally to 4,4-disubstituted-
3,4-dihydro-2(1H)-quinazolinones which are useful as
inhibitors of HIV reverse transcriptase, pharmaceutical
compositions and diagnostic kits comprising the same, methods
of using the same for treating viral infection or as assay
standards or reagents, and intermediates and processes for
making the same.

15

BACKGROUND OF THE INVENTION

Two distinct retroviruses, human immunodeficiency virus
(HIV) type-1 (HIV-1) or type-2 (HIV-2), have been
etiologically linked to the immunosuppressive disease,
acquired immunodeficiency syndrome (AIDS). HIV seropositive
individuals are initially asymptomatic but typically develop
AIDS related complex (ARC) followed by AIDS. Affected
individuals exhibit severe immunosuppression which
predisposes them to debilitating and ultimately fatal
opportunistic infections.

25

The disease AIDS is the end result of an HIV-1 or HIV-2
virus following its own complex life cycle. The virion life
cycle begins with the virion attaching itself to the host
human T-4 lymphocyte immune cell through the bonding of a
glycoprotein on the surface of the virion's protective coat
with the CD4 glycoprotein on the lymphocyte cell. Once
attached, the virion sheds its glycoprotein coat, penetrates
into the membrane of the host cell, and uncoats its RNA. The
virion enzyme, reverse transcriptase, directs the process of
transcribing the RNA into single-stranded DNA. The viral RNA
is degraded and a second DNA strand is created. The now
double-stranded DNA is integrated into the human cell's genes
and those genes are used for virus reproduction.

35

At this point, RNA polymerase transcribes the integrated DNA into viral RNA. The viral RNA is translated into the precursor gag-pol fusion polyprotein. The polyprotein is then cleaved by the HIV protease enzyme to yield the mature
5 viral proteins. Thus, HIV protease is responsible for regulating a cascade of cleavage events that lead to the virus particle's maturing into a virus that is capable of full infectivity.

The typical human immune system response, killing the
10 invading virion, is taxed because the virus infects and kills the immune system's T cells. In addition, viral reverse transcriptase, the enzyme used in making a new virion particle, is not very specific, and causes transcription mistakes that result in continually changed glycoproteins on
15 the surface of the viral protective coat. This lack of specificity decreases the immune system's effectiveness because antibodies specifically produced against one glycoprotein may be useless against another, hence reducing the number of antibodies available to fight the virus. The
20 virus continues to reproduce while the immune response system continues to weaken. Eventually, the HIV largely holds free reign over the body's immune system, allowing opportunistic infections to set in and without the administration of antiviral agents, immunomodulators, or both, death may
25 result.

There are at least three critical points in the virus's life cycle which have been identified as possible targets for antiviral drugs: (1) the initial attachment of the virion to the T-4 lymphocyte or macrophage site, (2) the transcription
30 of viral RNA to viral DNA (reverse transcriptase, RT), and (3) the processing of gag-pol protein by HIV protease.


Inhibition of the virus at the second critical point, the viral RNA to viral DNA transcription process, has provided a number of the current therapies used in treading
35 AIDS. This transcription must occur for the virion to reproduce because the virion's genes are encoded in RNA and the host cell reads only DNA. By introducing drugs that

block the reverse transcriptase from completing the formation of viral DNA, HIV-1 replication can be stopped.

A number of compounds that interfere with viral replication have been developed to treat AIDS. For example, nucleoside analogs, such as 3'-azido-3'-deoxythymidine (AZT), 2',3'-dideoxycytidine (ddC), 2',3'-dideoxythymidine (d4T), 2',3'-dideoxyinosine (ddI), and 2',3'-dideoxy-3'-thiacytidine (3TC) have been shown to be relatively effective in halting HIV replication at the reverse transcriptase (RT) stage.


An active area of research is in the discovery of non-nucleoside HIV reverse transcriptase inhibitors. As an example, it has been found that certain benzoxazinones and quinazolinones are active in the inhibition of HIV reverse transcriptase, the prevention or treatment of infection by HIV and the treatment of AIDS.

U.S. 5,519,021 describe reverse transcriptase inhibitors which are benzoxazinones of the formula:



wherein X is a halogen, Z may be O.

EP 0,530,994 and WO 93/04047 describe HIV reverse transcriptase inhibitors which are quinazolinones of the formula A:




A

wherein G is a variety of groups, R³ and R⁴ may be H, Z may be O, R² may be unsubstituted alkyl, unsubstituted alkenyl, unsubstituted alkynyl, unsubstituted cycloalkyl, unsubstituted heterocycle, and optionally substituted aryl, and R¹ may be a variety of groups including substituted alkyl.

WO 95/12583 also describes HIV reverse transcriptase inhibitors of formula A. In this publication, G is a variety of groups, R³ and R⁴ may be H, Z may be O, R² is substituted alkenyl or substituted alkynyl, and R¹ is cycloalkyl, alkynyl, alkenyl, or cyano. WO 95/13273 illustrates the asymmetric synthesis of one of the compounds of WO 95/12583, (S)-(-)-6-chloro-4-cyclopropyl-3,4-dihydro-4((2-pyridyl)ethynyl)-2(1H)-quinazolinone.

Synthetic procedures for making quinazolinones like those described above are detailed in the following references: Houpiis et al, *Tetr. Lett.* **1994**, 35(37), 6811-6814; Tucker et al, *J. Med. Chem.* **1994**, 37, 2437-2444; and, Huffman et al, *J. Org. Chem.* **1995**, 60, 1590-1594.

DE 4,320,347 illustrates quinazolinones of the formula:



wherein R is a phenyl, carbocyclic ring, or a heterocyclic ring. Compounds of this sort are not considered to be part of the present invention.

Even with the current success of reverse transcriptase inhibitors, it has been found that HIV patients can become resistant to a single inhibitor. Thus, it is desirable to develop additional inhibitors to further combat HIV infection.

SUMMARY OF THE INVENTION

Accordingly, one object of the present invention is to provide novel reverse transcriptase inhibitors.

It is another object of the present invention to provide a novel method for treating HIV infection which comprises administering to a host in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt form thereof.

It is another object of the present invention to provide a novel method for treating HIV infection which comprises


administering to a host in need thereof a therapeutically effective combination of (a) one of the compounds of the present invention and (b) one or more compounds selected from the group consisting of HIV reverse transcriptase inhibitors and HIV protease inhibitors.

It is another object of the present invention to provide pharmaceutical compositions with reverse transcriptase inhibiting activity comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt form thereof.

It is another object of the present invention to provide a method of inhibiting HIV present in a body fluid sample which comprises treating the body fluid sample with an effective amount of a compound of the present invention.

It is another object of the present invention to provide a kit or container containing at least one of the compounds of the present invention in an amount effective for use as a standard or reagent in a test or assay for determining the ability of a potential pharmaceutical to inhibit HIV reverse transcriptase, HIV growth, or both.


These and other objects, which will become apparent during the following detailed description, have been achieved by the inventors' discovery that compounds of formula (I):



wherein R¹, R², R³, and R⁸ are defined below, stereoisomeric forms, mixtures of stereoisomeric forms, or pharmaceutically acceptable salt forms thereof, are effective reverse transcriptase inhibitors.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[1] Thus, in a first embodiment, the present invention provides a novel compound of formula I:



I

or a stereoisomer or pharmaceutically acceptable salt
 5 thereof, wherein:

R¹ is C₁₋₃ alkyl substituted with 1-7 halogen;

10 R² is selected from C₁₋₅ alkyl substituted with 1-2 R⁴, C₂₋₅ alkenyl substituted with 1-2 R⁴, and C₂₋₅ alkynyl substituted with 1 R⁴;

15 R³, at each occurrence, is independently selected from C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, F, Cl, Br, I, NR⁵R^{5a}, NO₂, CN, C(O)R⁶, NHC(O)R⁷, and NHC(O)NR⁵R^{5a};

alternatively, if two R³'s are present and are attached to adjacent carbons, then they may combine to form -OCH₂O-;

20 R⁴ is selected from C₃₋₅ cycloalkyl substituted with 0-2 R³, phenyl substituted with 0-5 R³, and a 5-6 membered heterocyclic system containing 1-3 heteroatoms selected from O, N, and S, substituted with 0-2 R³;

25 R⁵ and R^{5a} are independently selected from H and C₁₋₃ alkyl;

R⁶ is selected from H, OH, C₁₋₄ alkyl, C₁₋₄ alkoxy, and NR⁵R^{5a};

R⁷ is selected from C₁₋₃ alkyl and C₁₋₃ alkoxy;

30

R⁸ is selected from H, C₃₋₅ cycloalkyl, and C₁₋₃ alkyl; and,

n is selected from 0, 1, 2, 3, and 4.

35

[2] In a preferred embodiment, the present invention provides a novel compound of formula I, wherein:

R¹ is C₁₋₃ alkyl substituted with 1-7 halogen;

5

R² is selected from C₁₋₅ alkyl substituted with 1 R⁴, C₂₋₅ alkenyl substituted with 1 R⁴, and C₂₋₅ alkynyl substituted with 1 R⁴;

10 R³, at each occurrence, is independently selected from C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, F, Cl, Br, I, NR⁵R^{5a}, NO₂, CN, C(O)R⁶, NHC(O)R⁷, and NHC(O)NR⁵R^{5a};

15 alternatively, if two R³'s are present and are attached to adjacent carbons, then they may combine to form -OCH₂O-;

20 R⁴ is selected from C₃₋₅ cycloalkyl substituted with 0-2 R³, phenyl substituted with 0-2 R³, and a 5-6 membered heterocyclic system containing 1-3 heteroatoms selected from O, N, and S, substituted with 0-1 R³;

R⁵ and R^{5a} are independently selected from H, CH₃ and C₂H₅;

25 R⁶ is selected from H, OH, CH₃, C₂H₅, OCH₃, OC₂H₅, and NR⁵R^{5a};

R⁷ is selected from CH₃, C₂H₅, OCH₃, and OC₂H₅;

R⁸ is selected from H, cyclopropyl, CH₃ and C₂H₅; and,

30 n is selected from 0, 1, 2, and 3.

[3] In a more preferred embodiment, the present invention provides a novel compound of formula I, wherein:

35

R¹ is selected from CF₃, and C₂F₅;

R² is selected from C₁₋₃ alkyl substituted with 1 R⁴, C₂₋₃ alkenyl substituted with 1 R⁴, and C₂₋₃ alkynyl substituted with 1 R⁴;

5 R³, at each occurrence, is independently selected from C₁₋₃ alkyl, OH, C₁₋₃ alkoxy, F, Cl, Br, I, NR⁵R^{5a}, NO₂, CN, C(O)R⁶, NHC(O)R⁷, and NHC(O)NR⁵R^{5a};

10 alternatively, if two R³'s are present and are attached to adjacent carbons, then they may combine to form -OCH₂O-;

R⁴ is selected from C₃₋₅ cycloalkyl substituted with 0-2 R³, phenyl substituted with 0-2 R³, and a 5-6 membered heterocyclic system containing 1-3 heteroatoms selected
15 from O, N, and S, substituted with 0-1 R³;

R⁵ and R^{5a} are independently selected from H, CH₃ and C₂H₅;

R⁶ is selected from H, OH, CH₃, C₂H₅, OCH₃, OC₂H₅, and NR⁵R^{5a};
20

R⁷ is selected from CH₃, C₂H₅, OCH₃, and OC₂H₅;

R⁸ is selected from H, CH₃ and C₂H₅; and,

25 n is selected from 0, 1, and 2.

[4] In an even more preferred embodiment, the present invention provides a novel compound of formula I, wherein:
30

R¹ is CF₃;

R² is selected from C₁₋₃ alkyl substituted with 1 R⁴, C₂₋₃ alkenyl substituted with 1 R⁴, and C₂₋₃ alkynyl
35 substituted with 1 R⁴;

R^3 , at each occurrence, is independently selected from C_{1-3} alkyl, OH, C_{1-3} alkoxy, F, Cl, NR^5R^{5a} , NO_2 , CN, $C(O)R^6$, $NHC(O)R^7$, and $NHC(O)NR^5R^{5a}$;

5 alternatively, if two R^3 's are present and are attached to adjacent carbons, then they may combine to form $-OCH_2O-$;

R^4 is selected from cyclopropyl substituted with 0-1 R^3 , phenyl substituted with 0-2 R^3 , and a 5-6 membered
10 heterocyclic system containing 1-3 heteroatoms selected from O, N, and S, substituted with 0-1 R^3 , wherein the heterocyclic system is selected from 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-furanyl, 3-furanyl, 2-thienyl, 3-thienyl, 2-oxazolyl, 2-thiazolyl, 4-isoxazolyl, and 2-
15 imidazolyl;

R^5 and R^{5a} are independently selected from H, CH_3 and C_2H_5 ;


R^6 is selected from H, OH, CH_3 , C_2H_5 , OCH_3 , OC_2H_5 , and NR^5R^{5a} ;
20

R^7 is selected from CH_3 , C_2H_5 , OCH_3 , and OC_2H_5 ;

R^8 is selected from H, CH_3 and C_2H_5 ; and,


25 n is selected from 1 and 2.

[5] In a further preferred embodiment, wherein the compound is of formula Ia



Ia.

[5] In a further preferred embodiment, wherein the compound
35 is of formula Ib:



Ib.

5 [7] In a further preferred embodiment, the compound of formula I is selected from:

- 10 (+/-)-6-Chloro-4-cyclopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (+/-)-6-Chloro-4-(2-pyridyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- 15 (+/-)-6-Chloro-4-phenylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (+/-)-4-Cyclopropylethynyl-6-methoxy-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- 20 (+/-)-6-Methoxy-4-(2-pyridyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (+/-)-6-Methoxy-4-phenylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- 25 (+/-)-4-Cyclopropylethynyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (+/-)-5,6-Difluoro-4-(2-pyridyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- 30 (+/-)-5,6-Difluoro-4-phenylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

(+/-)-4-Cyclopropylethynyl-6-fluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

5 (+/-)-6-Fluoro-4-(2-pyridyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

(+/-)-6-Fluoro-4-phenylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

10 (+/-)-6-Fluoro-4-(2'-2-pyridyl)ethyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

(+/-)-6-Fluoro-4-phenylethyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

15 (-)-6-Chloro-4-cyclopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

20 (+)-6-Chloro-4-cyclopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

(+)-4-Cyclopropylethynyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

25 (-)-4-Cyclopropylethynyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;


(+)-4-E-Cyclopropylethenyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone; and,

30 (-)-6-Chloro-4-E-cyclopropylethenyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

or a pharmaceutically acceptable salt thereof.

35

[8] In a second embodiment, the present invention provides a novel compound of formula II:



II

or a stereoisomer or pharmaceutically acceptable salt thereof, wherein:

5

R^2 is $C\equiv C-R^{4a}$;

R^3 is selected from C_{1-4} alkyl, OH, C_{1-4} alkoxy, F, Cl, Br, I, NR^5R^{5a} , NO_2 , CN, $C(O)R^6$, $NHC(O)R^7$, and $NHC(O)NR^5R^{5a}$;

10

R^{4a} is selected from methyl, ethyl, n-propyl, i-propyl, i-butyl, t-butyl, and i-pentyl;

R^5 and R^{5a} are independently selected from H and C_{1-3} alkyl;

15

R^6 is selected from H, OH, C_{1-4} alkyl, C_{1-4} alkoxy, and NR^5R^{5a} ;

R^7 is selected from C_{1-3} alkyl and C_{1-3} alkoxy;

20

R^8 is selected from H, C_{3-5} cycloalkyl, and C_{1-3} alkyl; and,

n is selected from 0, 1, 2, 3, and 4.

25

[9] In another preferred embodiment, the present invention provides a novel compound of formula II, wherein:

R^2 is $C\equiv C-R^{4a}$;

30

R^3 is selected from C_{1-4} alkyl, OH, C_{1-4} alkoxy, F, Cl, Br, I, NR^5R^{5a} , NO_2 , CN, $C(O)R^6$, and $NHC(O)R^7$;

R^{4a} is selected from methyl, ethyl, n-propyl, i-propyl, i-butyl, t-butyl, and i-pentyl;

35

R⁵ and R^{5a} are independently selected from H, CH₃ and C₂H₅;

R⁶ is selected from H, OH, CH₃, C₂H₅, OCH₃, OC₂H₅, and NR⁵R^{5a};


5 R⁷ is selected from CH₃, C₂H₅, OCH₃, and OC₂H₅;

R⁸ is selected from H, cyclopropyl, CH₃ and C₂H₅; and,

n is selected from 0, 1, and 2.

10


[10] In a further preferred embodiment, wherein the compound is of formula IIa



15

IIa.

[11] In a further preferred embodiment, wherein the compound is of formula IIb:



20

IIb.

[12] In another more preferred embodiment, the compound of
25 formula II is selected from:

(+/-)-6-Chloro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

30 (+/-)-6-Chloro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

- (+/-)-4-Isopropylethynyl-6-methoxy-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- 5 (+/-)-5,6-Difluoro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (+/-)-5,6-Difluoro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- 10 (+/-)-5,6-Difluoro-4-isopentyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (+/-)-6-Fluoro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- 15 (+/-)-6-Fluoro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (-)-5,6-Difluoro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- 20 (+)-5,6-Difluoro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (-)-5,6-Difluoro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone; and,
- 25 (+)-5,6-Difluoro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- 30 or a pharmaceutically acceptable salt thereof.

In a third embodiment, the present invention provides a
35 novel pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of formula I or II or pharmaceutically acceptable salt form thereof.

In a fourth embodiment, the present invention provides a novel method for treating HIV infection which comprises administering to a host in need of such treatment a therapeutically effective amount of a compound of formula I or II or pharmaceutically acceptable salt form thereof.

In a fifth embodiment, the present invention provides a novel method of treating HIV infection which comprises administering, in combination, to a host in need thereof a therapeutically effective amount of:

- (a) a compound of formula I or II; and,
- (b) at least one compound selected from the group consisting of HIV reverse transcriptase inhibitors and HIV protease inhibitors.

In another preferred embodiment, the reverse transcriptase inhibitor is selected from AZT, 3TC, ddI, ddC, d4T, delavirdine, TIBO derivatives, BI-RG-587, nevirapine, L-697,661, LY 73497, Ro 18,893, loviride, trovirdine, MKC-442, and HBY 097, and the protease inhibitor is selected from saquinavir, ritonavir, indinavir, VX-478, nelfinavir, KNI-272, CGP-61755, U-140690, and ABT-378.

In an even more preferred embodiment, the reverse transcriptase inhibitor is selected from AZT and 3TC and the protease inhibitor is selected from saquinavir, ritonavir, nelfinavir, and indinavir.

In a still further preferred embodiment, the reverse transcriptase inhibitor is AZT.

In another still further preferred embodiment, the protease inhibitor is indinavir.

- 5 In a sixth embodiment, the present invention provides a pharmaceutical kit useful for the treatment of HIV infection, which comprises a therapeutically effective amount of:
- (a) a compound of formula I or II; and,
 - (b) at least one compound selected from the group
- 10 consisting of HIV reverse transcriptase inhibitors and HIV protease inhibitors, in one or more sterile containers.

- 15 In a seventh embodiment, the present invention provides a novel method of inhibiting HIV present in a body fluid sample which comprises treating the body fluid sample with an effective amount of a compound of formula I or II.

- 20 In a eighth embodiment, the present invention to provides a novel a kit or container comprising a compound of formula I or II in an amount effective for use as a standard or reagent in a test or assay for determining the ability of a potential pharmaceutical to inhibit HIV reverse
- 25 transcriptase, HIV growth, or both.

DEFINITIONS

- As used herein, the following terms and expressions have the indicated meanings. It will be appreciated that the
- 30 compounds of the present invention contain an asymmetrically substituted carbon atom, and may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis, from optically active starting
- 35 materials. All chiral, diastereomeric, racemic forms and all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomer form is specifically indicated.

The processes of the present invention are contemplated to be practiced on at least a multigram scale, kilogram scale, multikilogram scale, or industrial scale. Multigram scale, as used herein, is preferably the scale wherein at least one starting material is present in 10 grams or more, more preferably at least 50 grams or more, even more preferably at least 100 grams or more. Multikilogram scale, as used herein, is intended to mean the scale wherein more than one kilogram of at least one starting material is used. Industrial scale as used herein is intended to mean a scale which is other than a laboratory scale and which is sufficient to supply product sufficient for either clinical tests or distribution to consumers.

As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. Examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, and s-pentyl. "Haloalkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogen (for example $-C_vF_w$ where $v = 1$ to 3 and $w = 1$ to $(2v+1)$). Examples of haloalkyl include, but are not limited to, trifluoromethyl, trichloromethyl, pentafluoroethyl, and pentachloroethyl. "Alkoxy" represents an alkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge. Examples of alkoxy include, but are not limited to, methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, s-butoxy, t-butoxy, n-pentoxy, and s-pentoxy. "Cycloalkyl" is intended to include saturated ring groups, such as cyclopropyl, cyclobutyl, or cyclopentyl. "Alkenyl" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain, such as ethenyl, propenyl and the like. "Alkynyl" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more triple carbon-carbon bonds which may occur in any stable

point along the chain, such as ethynyl, propynyl and the like.

"Halo" or "halogen" as used herein refers to fluoro, chloro, bromo and iodo. "Counterion" is used to represent a small, negatively charged species such as chloride, bromide, hydroxide, acetate, sulfate and the like.

As used herein, "aryl" or "aromatic residue" is intended to mean an aromatic moiety containing the specified number of carbon atoms, such as phenyl or naphthyl. As used herein, "carbocycle" or "carbocyclic residue" is intended to mean any stable 3- to 5- membered monocyclic ring, which may be saturated or partially unsaturated. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, biphenyl, naphthyl, indanyl, adamantyl, or tetrahydronaphthyl (tetralin).

As used herein, the term "heterocycle" or "heterocyclic system" is intended to mean a stable 5- to 6- membered monocyclic heterocyclic ring which is saturated partially unsaturated or unsaturated (aromatic), and which consists of carbon atoms and from 1 to 3 heteroatoms independently selected from the group consisting of N, O and S. The nitrogen and sulfur heteroatoms may optionally be oxidized. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. If specifically noted, a nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. As used herein, the term "aromatic heterocyclic system" is intended to mean a stable 5- to 6- membered monocyclic heterocyclic aromatic ring which consists of carbon atoms and from 1 to 3 heterotams independently selected from the group consisting of N, O and S. It is preferred that the total

number of S and O atoms in the aromatic heterocycle is not more than 1.

Examples of heterocycles include, but are not limited to, 2-pyrrolidonyl, 2H-pyrrolyl, 4-piperidonyl, 6H-1,2,5-thiadiazinyl, 2H,6H-1,5,2-dithiazinyl, furanyl, furazanyl, 5 imidazolidinyl, imidazoliny, imidazolyl, isoxazolyl, morpholinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl., oxazolyl, piperazinyl, piperidinyl, 10 pteridinyl, piperidonyl, 4-piperidonyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, pyrrolyl, tetrahydrofuranyl, 6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thiazolyl, thienyl, 15 thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, and 1,3,4-triazolyl. Preferred heterocycles include, but are not limited to, pyridinyl, furanyl, thienyl, 20 pyrrolyl, pyrazolyl, imidazolyl, and oxazolidinyl. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

As used herein, "HIV reverse transcriptase inhibitor" is intended to refer to both nucleoside and non-nucleoside 25 inhibitors of HIV reverse transcriptase (RT). Examples of nucleoside RT inhibitors include, but are not limited to, AZT, ddC, ddI, d4T, and 3TC. Examples of non-nucleoside RT inhibitors include, but are no limited to, delavirdine (Pharmacia and Upjohn U90152S), TIBO derivatives, BI-RG-587, 30 nevirapine (Boehringer Ingelheim), L-697,661, LY 73497, Ro 18,893 (Roche), loviride (Janssen), trovirdine (Lilly), MKC-442 (Triangle), and HBY 097 (Hoechst).

As used herein, "HIV protease inhibitor" is intended to refer to compounds which inhibit HIV protease. Examples 35 include, but are not limited, saquinavir (Roche, Ro31-8959), ritonavir (Abbott, ABT-538), indinavir (Merck, MK-639), VX-478 (Vertex/Glaxo Wellcome), nelfinavir (Agouron, AG-1343), KNI-272 (Japan Energy), CGP-61755 (Ciba-Geigy), U-140690

(Pharmacia and Upjohn), and ABT-378. Additional examples include the cyclic protease inhibitors disclosed in WO93/07128, WO 94/19329, WO 94/22840, and PCT Application Number US96/03426.

5 As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid
10 salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic
15 inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic,
20 stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

25 The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a
30 stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical*
35 *Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication commensurate with a reasonable benefit/risk ratio.

"Prodrugs" are intended to include any covalently bonded carriers which release the active parent drug according to formula (I) or other formulas or compounds of the present invention *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of the present invention, for example formula (I), are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of the present invention wherein the hydroxy or amino group is bonded to any group that, when the prodrug is administered to a mammalian subject, cleaves to form a free hydroxyl or free amino, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of the present invention, and the like.


"Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent. Only stable compounds are contemplated by the present invention.

"Substituted" is intended to indicate that one or more hydrogens on the atom indicated in the expression using "substituted" is replaced with a selection from the indicated group(s), provided that the indicated atom's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., =O) group, then 2 hydrogens on the atom are replaced.


"Therapeutically effective amount" is intended to include an amount of a compound of the present invention or an amount of the combination of compounds claimed effective to inhibit HIV infection or treat the symptoms of HIV infection in a host. The combination of compounds is preferably a synergistic combination. Synergy, as described for example by Chou and Talalay, Adv. Enzyme Regul. 22:27-55 (1984), occurs when the effect (in this case, inhibition of HIV replication) of the compounds when administered in combination is greater than the additive effect of the compounds when administered alone as a single agent. In general, a synergistic effect is most clearly demonstrated at suboptimal concentrations of the compounds. Synergy can be in terms of lower cytotoxicity, increased antiviral effect, or some other beneficial effect of the combination compared with the individual components.

SYNTHESIS

The compounds of the present invention can be prepared in a number of ways well known to one skilled in the art of organic synthesis. The compounds of the present invention can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art. Preferred methods include but are not limited to those methods described below. Each of the references cited below are hereby incorporated herein by reference.




Scheme 1 illustrates a method of preparing keto-anilines from an appropriately substituted 2-aminobenzoic acid. The acid is converted to its N-methoxy-N-methyl amide derivative which can then be displaced to obtain the R¹-substituted ketone. The keto-anilines are useful intermediates for the presently claimed compounds.




Scheme 2 describes another method of preparing keto-anilines, this time from an appropriately substituted aniline. After iodination and amine protection, a group such as trifluoromethyl can be introduced using a strong base and ethyl trifluoroacetate. Deprotection provides the keto-aniline. Additional means of preparing keto-anilines are known to one of skill in the art, e.g., Houpis et al, *Tetr. Lett.* **1994**, 35(37), 6811-6814, the contents of which are hereby incorporated herein by reference.

SCHEME 3



Another method of making 2-trifluoroacetylanilines is shown in Scheme 3. After forming the protected aniline, the amide is then reduced and the trifluoromethyl group added. Oxidation with an oxidant, such as MnO_2 , provides the useful intermediate.

SCHEME 4



Using the general method detailed in Scheme 4, one can prepare compounds of the present invention. Keto-aniline 1,

which may be prepared by the methods described in Schemes 1 and 2, is treated with trimethylsilyl isocyanate in dry tetrahydrofuran in the presence of dimethylaminopyridine followed by tetrabutylammonium fluoride to give the hydroxy-urea 2. The hydroxy-urea 2 is then dehydrated with a dehydrating agent such as 4Å molecular sieves in refluxing toluene or xylenes to give the ketimine 3. A substituted acetylenic R² group is added by treating the ketimine 3 with a lithium acetylide, which is prepared in a separate vessel by reacting the corresponding substituted acetylene with n-butyllithium in dry tetrahydrofuran, to give the 4,4-disubstituted 3,4-dihydro-2(1H)-quinazolinone 4, a compound of formula I. The acetylenic bond of the compound 4 may be reduced, e.g., by catalytic hydrogenation, to give the corresponding alkenyl group (not shown) or the saturated compound 5.

Other R² groups may also be introduced by directly reacting the imine 3 with a lithiate R²Li or a Grignard reagent R²MgX in the presence or absence of Lewis acid catalyst, such as BF₃ etherate. See also Huffman et al, *J. Org. Chem.* **1995**, *60*, 1590-1594, the contents of which are hereby incorporated herein by reference.

In certain instances, one enantiomer of a compound of Formula I or II may display superior activity compared with the other. When required, separation of the racemic material can be achieved by HPLC using a chiral column as exemplified in Examples 27-34 (Scheme 4) or by a resolution using a resolving agent such as camphonic chloride as in Thomas J. Tucker, et al, *J. Med. Chem.* **1994**, *37*, 2437-2444. A chiral compound of Formula I may also be directly synthesized using a chiral catalyst or a chiral ligand, e.g. Mark A. Huffman, et al, *J. Org. Chem.* **1995**, *60*, 1590-1594.


Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

Examples

Abbreviations used in the Examples are defined as follows: "°C" for degrees Celsius, "d" for doublet, "dd" for doublet of doublets, "eq" for equivalent or equivalents, "g" for gram or grams, "mg" for milligram or milligrams, "mL" for milliliter or milliliters, "H" for hydrogen or hydrogens, "hr" for hour or hours, "m" for multiplet, "M" for molar, "min" for minute or minutes, "MHz" for megahertz, "MS" for mass spectroscopy, "nmr" or "NMR" for nuclear magnetic resonance spectroscopy, "t" for triplet, "TLC" for thin layer chromatography, "EDAC" for 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, "DIPEA" for diisopropylethylamine, "TBAF" for tetrabutylammonium fluoride, "LAH" for lithium aluminium hydride, and "TEA" for triethylamine.

Example 1

Preparation of (+/-)-6-Chloro-4-cyclopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone ($R^4 =$ Cyclopropyl)



Step 1. Synthesis of II-a from I-a.

To a solution of compound I-a (4.55 g, 20.2 mmol) in anhydrous THF (40 mL) was added dimethylaminopyridine (0.25 g, 2.02 mmol) and trimethylsilyl isocyanate (6.05 g, 7.11 mL, 52.5 mmol). The mixture was stirred at room temperature for

approximately 16 hours, then tetrabutylammonium fluoride (21 mL of 1 M solution in THF) was added. The thick slurry was diluted with additional THF (20 mL) and stirred at room temperature for 0.5 hours. The THF was removed under reduced pressure, the residue was taken up in EtOAc (100 mL) and washed sequentially with 1 N HCl (70 mL), saturated aqueous NaHCO₃ (70 mL) and saturated aqueous NaCl (50 mL). The organic phase was dried over MgSO₄, filtered and concentrated under reduced pressure to afford a light yellow solid. The yellow color was removed upon trituration with hexanes to afford IIa (5.09 g, 94%) as a white solid: ¹H NMR (300 MHz, acetone-d₆) δ 9.06 (br s, 1 H), 7.48 (s, 1 H), 7.40 (br s, 1 H), 7.34 (dd, J = 8.8, 2.6 Hz, 1 H), 6.97 (d, J = 8.8 Hz, 1 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -86.33, -86.35; IR (KBr Pellet) 1724, 1678, 1398, 1198, 1174 cm⁻¹; MS (CI) m/e 266 (MH⁺, 100).

Step 2. Synthesis of III-a from II-a.

A suspension of II-a (5.09 g, 19.1 mmol) in toluene (150 mL) containing 4 Å molecular sieves (approximately 100 mg) was heated at reflux for 16 hours. The resulting clear yellow solution was cooled to room temperature, the precipitated solids were dissolved in acetone and the molecular sieves were removed by vacuum filtration. The filtrate was concentrated under reduced pressure, and triturated with hexanes to afford III-a (4.25 g, 89%) as a yellow solid: ¹H NMR (300 MHz, acetone-d₆) δ 7.86-7.82 (m, 2 H), 7.61 (d, J = 8.8 Hz, 1 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -67.88.

Step 3. Synthesis of IV-a from IIIa.

A solution of cyclopropylacetylene (13.0 mL of 30 wt% solution in toluene/THF/hexanes, 59.0 mmol) in anhydrous THF (118 mL) was cooled to -78 °C, treated with n-BuLi (32.8 mL of 1.6 M solution in hexanes, 52.4 mmol), warmed to 0 °C in an ice bath, and aged for 0.5 h. To a solution of III-a

(3.12 g, 12.6 mmol) in anhydrous THF (66 mL) at -78 °C was added the lithium acetylide over approximately 10 minutes. To this was added boron trifluoride etherate (0.89 g, 0.80 mL, 6.28 mmol), followed by removal of the cooling bath. The reaction was allowed to reach room temperature and stirred at room temperature for 4 hours before quenching with 1 M citric acid (100 mL). The mixture was concentrated under reduced pressure to 1/2 original volume, diluted with EtOAc (200 mL), the aqueous phase was removed and the organic phase was sequentially washed with saturated aqueous NaHCO₃ (100 mL), and saturated aqueous NaCl (100 mL). The organic phase was dried over MgSO₄, filtered and concentrated under reduced pressure. The crude material was purified by flash chromatography (3% MeOH/CH₂Cl₂) to afford a thick yellow oil from which was obtained crystalline IV-a (R⁴ = cyclopropyl) (3.85 g, 97%) as a white solid: mp 86.6-88 °C; ¹H NMR (300 MHz, acetone-d₆) δ 8.95 (br s, 1 H), 7.51 (br s, 1 H), 7.43 (br s, 1 H), 7.40 (dd, J = 8.8, 2.4 Hz, 1 H), 7.02 (d, J = 8.8 Hz, 1 H), 1.49-1.41 (m, 1 H), 0.93-0.82 (m, 1 H), 0.77-0.74 (m, 1 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -82.96; IR (KBr Pellet) 1696, 1172 cm⁻¹; MS (CI) m/e calc'd for C₁₄H₁₀ClF₃N₂O: 315.051201, found 315.051626; 315 (MH⁺, 51), 332 (M+NH₄⁺, 100); Analysis calc'd for C₁₄H₁₀N₂ClF₃O·0.25 H₂O: C, 52.68; H, 3.32; N, 8.78; found: C, 52.61; H, 3.35; N, 8.28.

Example 2

Preparation of (+/-)-6-Chloro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Isopropyl)

A solution of III-a (50 mg, 0.201 mmol) was treated with the lithium acetylide derived from 3-methyl-1-butyne (62 mg, 93 mL, 0.905 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by flash chromatography (35% EtOAc/hexanes) to afford 26mg (41%) of the desired product: ¹H NMR (300 MHz,) δ 9.08 (br s, 1 H), 7.59 (br s, 1 H), 7.53 (br s, 1 H), 7.40 (dd, J = 8.4,

2.2 Hz, 1 H), 7.02 (d, $J = 8.8$ Hz, 1 H), 2.81-2.68 (m, 1 H), 1.20 (dd, $J = 6.6$ Hz, 6H); ^{19}F NMR (282 MHz, acetone- d_6) δ -83.05; MS (CI) m/e calc'd for $\text{C}_{14}\text{H}_{12}\text{ClF}_3\text{N}_2\text{O}$: 317.066851, found 317.069433; 317 (MH^+ , 43), 334 ($\text{M}+\text{NH}_4^+$, 100).

5

Example 3

Preparation of (+/-)-6-Chloro-4-(2-pyridyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone ($\text{R}^4 =$ 2-Pyridyl)

10

A solution of III-a (100 mg, 0.402 mmol) was treated with the the lithium acetylide derived from 2-ethynylpyridine (0.19 g, 1.81 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by HPLC (2.5% MeOH/ CH_2Cl_2) to afford 85 mg (60%) of the desired product: mp 105 °C dec.; ^1H NMR (300 MHz, acetone- d_6) δ 9.14 (br s, 1 H), 8.64-8.61 (m, 1 H), 7.89-7.84 (m, 2 H), 7.70-7.66 (m, 2 H), 7.48-7.43 (m, 2 H), 7.09 (d, $J = 8.8$ Hz, 1 H); ^{19}F NMR (282 MHz, acetone- d_6) δ -82.48; IR (KBr Pellet) 1704, 1430, 1186 cm^{-1} ; MS (CI) m/e calc'd for $\text{C}_{16}\text{H}_{10}\text{ClF}_3\text{N}_3\text{O}$: 352.046450, found 352.046956; 352 (MH^+ , 100); Analysis calc'd for $\text{C}_{16}\text{H}_9\text{ClF}_3\text{N}_3\text{O} \cdot 0.125 \text{ H}_2\text{O}$: C, 54.3; H, 2.56; N, 11.9; found: C, 54.71; H, 3.03; N, 11.3.

25

Example 4

Preparation of (+/-)-6-Chloro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone ($\text{R}^4 =$ Ethyl)

A solution of III-a (100 mg, 0.402 mmol) was treated with the the lithium acetylide derived from 1-butyne (109 mg, 2.01 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by HPLC (2.5% MeOH/ CH_2Cl_2) to afford 79 mg (65%) of the desired product: ^1H NMR (300 MHz, acetone- d_6) δ 9.05 (br s, 1 H), 7.54 (br s, 2 H), 7.41-7.39 (m, 1 H), 7.02 (d, $J = 8.4$ Hz, 1 H), 2.36-2.32 (m, 2 H), 2.18-1.13 (m, 3 H); ^{19}F NMR (282 MHz, acetone- d_6) δ

-82.99; MS (CI) m/e calc'd for $C_{13}H_{10}ClF_3N_2O$: 303.051201, found 303.051882; 303 (MH^+ , 55), 320 ($M+NH_4^+$, 100).


Example 5

5 Preparation of (+/-)-6-Chloro-4-phenylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R^4 = Phenyl)

A solution of III-a (100 mg, 0.402 mmol) was treated with the lithium acetylide derived from phenylacetylene (185 mg, 1.81 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by HPLC (2.5% MeOH/ CH_2Cl_2) to afford 54 mg (38%) of the desired product: 1H NMR (300 MHz, acetone- d_6) δ 9.07 (br s, 1 H), 7.74 (br s, 1 H), 7.67 (br s, 1 H), 7.62-7.58 (m, 2 H), 7.48-7.40 (m, 4H), 7.08 (d, J = 8.4 Hz, 1 H); ^{19}F NMR (282 MHz, acetone- d_6) δ -82.67; IR (KBr Pellet) 1696, 1186 cm^{-1} ; MS (CI) m/e calc'd for $C_{17}H_{11}ClF_3N_2O$: 351.051201, found 351.051704; 351 (MH^+ , 51), 368 ($M+NH_4^+$, 100); Analysis calc'd for $C_{17}H_{10}ClF_3N_2O \cdot 0.25 H_2O$: C, 57.48; H, 2.98; N, 7.89; found: C, 57.00; H, 3.03; N, 7.48.

Example 6

25 Preparation of (+/-)-4-Cyclopropylethynyl-6-methoxy-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R^4 = Cyclopropyl)



30 Step 1. Synthesis of VI-a from V-a.

A solution of V-a (0.50 g, 2.28 mmol) was treated with dimethylaminopyridine and trimethylsilyl isocyanate as described in Step 1 of Example 1 to afford 0.58 g (97%) of the desired product: ^1H NMR (300 MHz, acetone- d_6) δ 8.81 (br s, 1 H), 7.17 (br s, 1 H), 7.11 (br s, 1 H), 7.00-6.92 (m, 2 H), 6.83 (s, 1 H), 3.76 (s, 3 H); ^{19}F NMR (282 MHz, acetone- d_6) δ -85.99.

Step 2. Synthesis of VII-a from VI-a.

A solution of VI-a (0.58 g, 2.21 mmol) was heated in toluene at reflux as described in Step 2 of Example 1 to afford 0.50 g (93%) of the desired product: ^1H NMR (300 MHz, acetone- d_6) δ 7.52 (br s, 2 H), 7.27 (s, 1 H), 3.90 (s, 3 H); ^{19}F NMR (282 MHz, acetone- d_6) δ -68.08.

Step 3. Synthesis of VIII-a from VII-a.

A solution of VII-a (100 mg, 0.410 mmol) was treated with the lithium acetylide derived from cyclopropylacetylene (0.41 mL of 30 wt% solution in toluene/THF/hexanes, 1.85 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by HPLC (2.5% MeOH/ CH_2Cl_2) to afford 103 mg (81%) of the desired product: ^1H NMR (300 MHz, acetone- d_6) δ 8.77 (br s, 1 H), 7.29 (br s, 1 H), 7.06 (br s, 1 H), 6.99-6.90 (m, 2 H), 3.77 (s, 3 H), 1.46-1.38 (m, 1 H), 0.91-0.85 (m, 2 H), 0.79-0.72 (m, 2 H); ^{19}F NMR (282 MHz, acetone- d_6) δ -82.61; MS (CI) m/e calc'd for $\text{C}_{15}\text{H}_{14}\text{F}_3\text{N}_2\text{O}_2$: 311.100738, found 311.099970; 311 (MH^+ , 100).

Example 7

Preparation of (+/-)-4-Isopropylethynyl-6-methoxy-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Isopropyl)

5
A solution of VII-a (100 mg, 0.410 mmol) was treated with the the lithium acetylide derived from 3-methyl-1-butyne (126 mg, 0.19 mL, 1.85 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was
10 purified by flash chromatography (2.5% MeOH/CH₂Cl₂) to afford 30 mg (24%) of the desired product: mp 228-229 °C; ¹H NMR (300 MHz, acetone-d₆) δ 8.72 (br s, 1 H), 7.27 (br s, 1 H), 7.10 (br s, 1 H), 7.00-6.91 (m, 2 H), 3.77 (s, 3 H), 2.73-2.67 (m, 1 H), 1.20 (dd, J = 7.0, 1.5 Hz, 6 H); ¹⁹F NMR (282
15 MHz, acetone-d₆) δ -82.71; IR (KBr Pellet) 1696, 1428, 1190, 1176 cm⁻¹; MS (CI) m/e calc'd for C₁₅H₁₆F₃N₂O₂: 313.116388, found 313.115871; 313 (MH⁺, 100), 330 (M+NH₄⁺, 15); Analysis calc'd for C₁₅H₁₅F₃N₂O₂: C, 57.69; H, 4.84; N, 8.97; found: C, 57.74; H, 5.01; N, 8.57.

20

Example 8

Preparation of (+/-)-6-Methoxy-4-(2-pyridyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = 2-Pyridyl)

25

A solution of VII-a (100 mg, 0.410 mmol) was treated with the the lithium acetylide derived from 2-ethynylpyridine (0.19 g, 1.85 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by
30 flash chromatography (2.5% MeOH/CH₂Cl₂) to afford 56 mg (39%) of the desired product: ¹H NMR (300 MHz, acetone-d₆) δ 8.81 (br s, 1 H), 8.61 (d, J = 4.8 Hz, 1 H), 7.88-7.82 (m, 1 H), 7.66 (d, J = 7.7 Hz, 1 H), 7.61 (br s, 1 H), 7.46-7.42 (m, 1 H), 7.23 (br s, 1 H), 7.06-6.97 (m, 2 H), 3.79 (s, 3 H); ¹⁹F
35 NMR (282 MHz, acetone-d₆) δ -82.13; IR (KBr Pellet) 1698, 1518, 1464, 1430, 1244, 1208, 1184 cm⁻¹; MS (CI) m/e calc'd for C₁₇H₁₃F₃N₃O₂: 348.095987, found 348.095629; 348 (MH⁺,

100); Analysis calc'd for $C_{17}H_{12}F_3N_3O_2 \cdot 0.25 C_3H_6O$: C, 58.92; H, 3.76; N, 11.61; found: C, 59.38; H, 4.04; N, 11.35.


Example 9

5 **Preparation of (+/-)-6-Methoxy-4-phenylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Phenyl)**

A solution of VII-a (100 mg, 0.410 mmol) was treated
10 with the the lithium acetylide derived from phenylacetylene
(0.19 g, 1.85 mmol) according to the procedure of Step 3 of
Example 1. The resulting crude material was purified by
flash chromatography (2.5% MeOH/CH₂Cl₂) to afford 34 mg (24%)
of the desired product: mp 206.2-207.7 °C; ¹H NMR (300 MHz,
15 acetone-d₆) δ 8.85 (br s, 1 H), 7.60-7.57 (m, 3 H), 7.49-7.39
(m, 3 H), 7.21 (br s, 1 H), 7.05-6.96 (m, 2H), 3.79 (s, 3 H);
¹⁹F NMR (282 MHz, acetone-d₆) δ -82.32; IR (KBr Pellet) 1696,
1516, 1430, 1236, 1204, 1184, 1128 cm⁻¹; MS (CI) m/e calc'd
for C₁₈H₁₄F₃N₂O₂: 347.100738, found 347.101482; 347 (MH⁺,
20 100), 364 (M+NH₄⁺, 48); Analysis calc'd for C₁₈H₁₃F₃N₂O₂: C,
62.43; H, 3.78; N, 8.10; found: C, 62.35; H, 3.58; N,
7.83.

Example 10**Preparation of (+/-)-4-Cyclopropylethynyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R^4 = Cyclopropyl)**

5



Step 1. Synthesis of X-a from IX-a.

10 A solution of IX-a (6.46 g, 28.7 mmol) was treated with dimethylaminopyridine and trimethylsilyl isocyanate as described in Step 1 of Example 1 to afford 6.74 g (88%) of the desired product: ¹H NMR (300 MHz, acetone-d₆) δ 9.13 (br s, 1 H), 7.45-7.32 (m, 2 H), 7.18 (br s, 1 H), 6.85-6.80 (m, 1 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -86.6 (d, 17.2, 3), -137.52-137.68 (m, 1), -148.47-148.59 (m, 1).

Step 2. Synthesis of XI-a from X-a.

20 A solution of X-a (6.74 g, 25.1 mmol) was heated in xylenes at reflux as described in Step 2 of Example 1, substituting xylenes for toluene, to afford 6.3 g (100%) of the desired product: ¹H NMR (300 MHz, acetone-d₆) δ 7.92-7.83 (m, 1 H), 7.46-7.44 (m, 1 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -70.7 (d, 38.7, 3), -136.72 (s, 1), -146.47-146.57 (m, 1).

Step 3. Synthesis of XII-a from XI-a.

A solution of XI-a (6.28 g, 25.1 mmol) was treated with the the lithium acetylide derived from cyclopropylacetylene (24.9 mL of 30 wt% solution in toluene/THF/hexanes, 0.113 mol) according to the procedure of Step 3 of Example 1. The resulting crude yellow oil was dissolved in acetone and concentrated under reduced pressure to deliver a yellow solid. Crystallization from acetone afforded 5.98 g (75%) of the desired material: mp 86.5-88.5 °C; ¹H NMR (300 MHz, acetone-d₆) δ 9.01 (br s, 1 H), 7.46 (br s, 1 H), 7.44-7.35 (m, 1 H), 6.86-6.81 (m, 1 H), 1.41-1.37 (m, 1 H), 0.90-0.83 (m, 1 H), 0.74-0.69 (m, 1 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -83.3 (d, J = 12.9, 1), -136.04-136.23 (m, 1), -148.14-148.26 (m, 1); IR (KBr Pellet) 1706, 1516, 1442, 1246, 1214, 1196 cm⁻¹; MS (CI) m/e calc'd for C₁₄H₁₀F₅N₂O: 317.071329, found 317.070836; 317 (MH⁺, 100), 334 (M+NH₄⁺, 62); Analysis calc'd for C₁₄H₉F₅N₂O: C, 53.17; H, 2.88; N, 8.87; found: C, 53.30; H, 3.16; N, 8.53.

20

Example 11

Preparation of (+/-)-5,6-Difluoro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Isopropyl)

25

A solution of XI-a (7.24 g, 28.9 mmol) was treated with the the lithium acetylide derived from 3-methyl-1-butyne (8.87 g, 13.3 mL, 0.130 mol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by flash chromatography (2.5% MeOH/CH₂Cl₂) to afford a yellow oil. Crystallization from acetone afforded 6.77 g (74%) of the desired product: mp 79-80 °C; ¹H NMR (300 MHz, acetone-d₆) δ 9.02 (br s, 1 H), 7.50 (br s, 1 H), 7.44-7.35 (m, 1 H), 6.87-6.82 (m, 1 H), 2.69-2.65 (m, 1 H), 1.17 (d, J = 7.0 Hz, 6H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -83.4 (d, J = 12.9, 1), -135.79-135.94 (m, 1), -148.14-148.26 (m, 1); MS (CI) m/e calc'd for C₁₄H₁₂F₅N₂O: 319.086979, found 319.087376; 319 (MH⁺, 100), 336 (M+NH₄⁺, 76).

Example 12**Preparation of (+/-)-5,6-Difluoro-4-(2-pyridyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = 2-Pyridyl)**

5

A solution of XI-a (100 mg, 0.400 mmol) was treated with the the lithium acetylide derived from 2-ethynylpyridine (0.19 g, 1.80 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by flash chromatography (4% MeOH/CH₂Cl₂) to afford 83 mg (59%) of the desired product: mp 219-220 °C; ¹H NMR (300 MHz, acetone-d₆) δ 9.15 (br s, 1 H), 8.61 (d, J = 4.4 Hz, 1 H), 7.88-7.82 (m, 2H), 7.63 (dd, J = 7.0, 1.1 Hz, 1 H), 7.47-7.42 (m, 2H), 6.94-6.88 (m, 1 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -82.8 (d, J = 12.9, 3), -135.78-135.93 (m, 1), -147.86-147.98 (m, 1); IR (KBr Pellet) 1712, 1470, 1450, 1430, 1416, 1264, 1238, 1226, 1198, 1186 cm⁻¹; MS (CI) m/e calc'd for C₁₆H₉F₅N₃O: 354.066578, found 354.067821; 354 (MH⁺, 100).

20

Example 13**Preparation of (+/-)-5,6-Difluoro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = 2-Ethyl)**

A solution of XI-a (100 mg, 0.400 mmol) was treated with the the lithium acetylide derived from 1-butyne (97 mg, 1.80 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by HPLC (2.5% MeOH/CH₂Cl₂) to afford 69 mg (57%) of the desired product: mp 191-194 °C; ¹H NMR (300 MHz, acetone-d₆) δ 9.03 (br s, 1 H), 7.50 (br s, 1 H), 7.45-7.35 (m, 1 H), 6.87-6.82 (m, 1 H), 2.34-2.27 (m, 2H), 1.20-1.15 (m, 3 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -83.3 (d, J = 12.9, 3), -135.79-135.98 (m, 1), -148.16-148.29 (m, 1); IR (KBr Pellet) 1704, 1686, 1518, 1444, 1244, 1210, 1192, 1172 cm⁻¹; MS (CI) m/e calc'd for C₁₃H₁₀F₅N₂O: 305.071329, found 305.071146; 305 (MH⁺, 100); Analysis calc'd for C₁₃H₉F₅N₂O: C, 51.33; H, 2.98; N, 9.22; found: C, 51.00; H, 2.79; N, 8.99.

Example 14

Preparation of (+/-)-5,6-Difluoro-4-phenylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Phenyl)

A solution of XI-a (100 mg, 0.400 mmol) was treated with the the lithium acetylide derived from phenylacetylene (0.18 g, 0.20 mL, 1.80 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by HPLC (2.5% MeOH/CH₂Cl₂) to afford 92 mg (65%) of the desired product: ¹H NMR (300 MHz, acetone-d₆) δ 9.14 (br s, 1 H), 7.80 (br s, 1 H), 7.57-7.54 (m, 2H), 7.49-7.40 (m, 4H), 6.92-6.87 (m, 1 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -83.0 (d, J = 12.9, 3), -136.08-136.27 (m, 1), -147.87-148.00 (m, 1); MS (CI) m/e calc'd for C₁₇H₁₀F₅N₂O: 353.071329, found 353.071716; 353 (MH⁺, 42), 370 (M+NH₄⁺, 100).

Example 15

Preparation of (+/-)-5,6-Difluoro-4-isopentyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Isopropyl)

Synthesis of XIII-a from XII-a.

A solution of XIII-a (R⁴ = isopropyl) (26 mg, 82 mmol) in ethanol (1 mL) and EtOAc (0.5 mL) was treated with 10% Pd on carbon (35 mg) under H₂ (1 atm) for 16 hours. The catalyst was removed by vacuum filtration through Celite and the filter cake was washed with EtOAc. The combined filtrates were concentrated under reduced pressure to afford 26 mg (100%) of the desired material. No further purification was necessary: ¹H NMR (300 MHz, acetone-d₆) δ 8.88 (br s, 1 H), 7.41-7.31 (m, 1 H), 6.89-6.81 (m, 2H), 2.55-2.50 (m, 1 H), 1.64-1.45 (m, 2H), 1.06-1.02 (m, 1 H), 0.89 (dd, J = 6.6, 2.2 Hz, 6H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -83.22 (d, J = 12.1, 3), -138.97-139.13 (m, 1), -148.46-148.58 (m, 1); IR (KBr Pellet) 1700, 1678, 1518, 1438, 1252, 1188, 1172 cm⁻¹; MS

(CI) m/e calc'd for $C_{14}H_{16}F_5N_2O$: 323.118280, found 323.116703; 323 (MH^+ , 100), 340 ($M+NH_4^+$, 57).

Example 16

5 **Preparation of (+/-)-4-Butyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R^4 = Ethyl)**


10 A solution of XIII-a (R^4 = ethyl) (20 mg, 66 μ mol) in ethanol (1 mL) and EtOAc (0.5 mL) was treated with 10% Pd on carbon under H_2 according to the procedure of Example 15. Purification by HPLC (2.5% MeOH/ CH_2Cl_2) afforded 12 mg (56%) of the desired product: 1H NMR (300 MHz, acetone- d_6) δ 8.89 (br s, 1 H), 7.41-7.32 (m, 1 H), 6.86-6.81 (m, 2H), 2.57-2.47 (m, 1 H), 1.56-1.15 (m, 5H), 0.88 (t, J = 7.3 Hz, 3 H); ^{19}F NMR (282 MHz, acetone- d_6) δ -83.19-83.24 (m, 1), -139.14 (s, 1), -148.49-148.62 (m, 1); MS (CI) m/e calc'd for $C_{13}H_{14}F_5N_2O$: 309.102629, found 309.103555; 309 (MH^+ , 100), 326 ($M+NH_4^+$, 62).

20

Example 17

Preparation of (+/-)-4-Cyclopropylethynyl-6-fluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Cyclopropyl)

5



Step 1. Synthesis of XV-a from XIV-a.

10 A solution of XIII-a (3.07 g, 14.8 mmol) was treated with dimethylaminopyridine and trimethylsilyl isocyanate as described in Step 1 of Example 1 to afford 2.81 g (76%) of the desired product.

15 Step 2. Synthesis of XVI-a from XV-a.

A solution of XV-a (6.74 g, 25.1 mmol) was heated in toluene at reflux as described in Step 2 of Example 1 to afford 0.73 g (94%) of the desired product.

20

Step 3. Synthesis of XVII-a from XVI-a.

A solution of XVI-a (100 mg, 0.431 mmol) was treated with the the lithium acetylide derived from cyclopropylacetylene (1.43 mL of 30 wt% solution in toluene/THF/hexanes, 1.94 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by HPLC (2.5% MeOH/CH₂Cl₂) to afford 44 mg (34%) of the desired product: mp 155 °C; ¹H NMR (300 MHz, acetone-d₆) δ 8.86 (br s, 1 H), 7.36 (br s, 1 H), 7.30-7.27 (m, 1 H), 7.22-7.15 (m, 1 H), 7.04-6.99 (m, 1 H), 1.47-1.42 (m, 1 H), 0.90-0.87 (m, 2 H), 0.76-0.75 (m, 2 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -82.86, -123.36-123.44; MS (CI) m/e calc'd for C₁₄H₁₁F₄N₂O: 299.080751, found 299.079976; 299 (MH⁺, 100).

Example 18

Preparation of (+/-)-6-Fluoro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Isopropyl)

A solution of XVI-a (100 mg, 0.431 mmol) was treated with the the lithium acetylide derived from 3-methyl-1-butyne (0.13 g, 0.20 mL, 1.94 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by HPLC (2.5% MeOH/CH₂Cl₂) to afford 24 mg (18%) of the desired product: mp 158 °C; ¹H NMR (300 MHz, acetone-d₆) δ 9.07 (br s, 1 H), 7.60 (br s, 1 H), 7.32-7.30 (m, 1 H), 7.24-7.16 (m, 1 H), 7.05-6.99 (m, 1 H), 2.77-2.67 (m, 1 H), 1.20 (dd, J = 7.0, 2.6 Hz, 6H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -82.95, -123.41-123.49; MS (301) m/e calc'd for C₁₄H₁₃F₄N₂O: 301.096401, found 301.096235; 301 (MH⁺, 100).

Example 19

Preparation of (+/-)-6-Fluoro-4-(2-pyridyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = 2-Pyridyl)

A solution of XVI-a (100 mg, 0.431 mmol) was treated with the the lithium acetylide derived from 2-ethynylpyridine (0.20 g, 1.94 mmol) according to the procedure of Step 3 of

Example 1. The resulting crude material was purified by HPLC (2.5% MeOH/CH₂Cl₂) to afford 65 mg (45%) of the desired product: mp 155 °C; ¹H NMR (300 MHz, acetone-d₆) δ 9.02 (br s, 1 H), 8.60 (d, J = 4.0 Hz, 1 H), 7.87-7.78 (m, 2 H), 7.66 (d, J = 7.7 Hz, 1 H), 7.45-7.41 (m, 2 H), 7.26-7.20 (m, 1 H), 7.09-7.05 (m, 1 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -82.36, -122.94-123.02; MS (CI) m/e calc'd for C₁₆H₁₀F₄N₃O: 336.076000, found 336.074156; 336 (MH⁺, 25).

10

Example 20**Preparation of (+/-)-6-Fluoro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Ethyl)**

15

A solution of XVI-a (100 mg, 0.431 mmol) was treated with the lithium acetylide derived from 1-butyne (0.10 g, 1.94 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by HPLC (2.5% MeOH/CH₂Cl₂) to afford 40 mg (33%) of the desired product: mp 190 °C; ¹H NMR (300 MHz, acetone-d₆) δ 8.86 (br s, 1 H), 7.38 (br s, 1 H), 7.34-7.31 (m, 1 H), 7.22-7.16 (m, 1 H), 7.05-7.00 (m, 1 H), 2.04-2.01 (m, 2 H), 1.19-1.14 (m, 3 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -75.392, -123.42-123.50; MS (CI) m/e calc'd for C₁₃H₁₁F₄N₂O: 287.080751, found 287.080740; 287 (MH⁺, 100).

25

Example 21**Preparation of (+/-)-6-Fluoro-4-phenylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Phenyl)**

30

A solution of XVI-a (100 mg, 0.431 mmol) was treated with the lithium acetylide derived from phenylacetylene (0.20 g, 0.21 mL, 1.94 mmol) according to the procedure of Step 3 of Example 1. The resulting crude material was purified by HPLC (2.5% MeOH/CH₂Cl₂) to afford 41 mg (28%) of the desired product: mp 107 °C; ¹H NMR (300 MHz, acetone-d₆) δ 9.00 (br s, 1 H), 7.69 (br s, 1 H), 7.63-7.59 (m, 2 H),

35

7.50-7.40 (m, 4H), 7.27-7.20 (m, 1 H), 7.10-7.05 (m, 1 H);
¹⁹F NMR (282 MHz, acetone-d₆) δ -82.56, -122.99-123.07; MS
(CI) m/e calc'd for C₁₇H₁₁F₄N₂O: 335.080751, found
335.082057; 335 (MH⁺, 74), 352 (M+NH₄⁺, 100).

5

Example 22

Preparation of (+/-)-6-Fluoro-4-isopentyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Isopropyl)

10

Synthesis of XVIII-a from XVII-a.

A solution of XVII-a (R⁴ = isopropyl) (26 mg, 87 mmol) in ethanol (1 mL) and EtOAc (0.5 mL) was treated with 10% Pd on carbon under H₂ according to the procedure of Example 15 to afford 15 mg (58%) of the desired product. No further purification was necessary: mp 179 °C; ¹H NMR (300 MHz, acetone-d₆) δ 7.02-6.97 (m, 2 H), 6.80-6.76 (m, 1 H), 2.18-2.09 (m, 2 H), 1.92-1.82 (m, 2 H), 1.52-1.45 (m, 1 H), 0.88-0.79 (m, 6 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -82.60, -123.72-123.84; MS (CI) m/e calc'd for C₁₄H₁₇F₄N₂O: 305.127707, found 305.126790; 305 (MH⁺, 100).

20

Example 23

Preparation of (+/-)-6-Fluoro-4-(2'-2-pyridyl)ethyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = 2-Pyridyl)

25

A solution of XVII-a (R⁴ = 2-pyridyl) (33 mg, 99 mmol) in ethanol (1 mL) and EtOAc (0.5 mL) was treated with 10% Pd on carbon under H₂ according to the procedure of Example 15 to afford 10 mg (30%) of the desired product. No further purification was necessary: mp 88 °C; ¹H NMR (300 MHz, acetone-d₆) δ 8.35 (d, J = 4.4 Hz, 1 H), 7.63 (dt, J = 7.7, 1.5 Hz, 1 H), 7.20-7.13 (m, 3 H), 7.04-6.98 (m, 1 H), 6.83-6.79 (m, 1 H), 2.84-2.78 (m, 1 H), 2.68-2.48 (m, 2 H), 2.27-2.06 (m, 1 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -82.58,

35

-123.26-123.34; MS (CI) *m/e* calc'd for C₁₆H₁₄F₄N₃O:
340.107300, found 340.107719; 340 (MH⁺, 100).

Example 24

5 **Preparation of (+/-)-4-Butyl-6-fluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Ethyl)**

10 A solution of XVII-a (R⁴ = ethyl) (24 mg, 84 mmol) in ethanol (1 mL) and EtOAc (0.5 mL) was treated with 10% Pd on carbon under H₂ according to the procedure of Example 15 to afford 24 mg (100%) of the desired product. No further purification was necessary: mp 198 °C; ¹H NMR (300 MHz, acetone-d₆) δ 7.03-6.97 (m, 2 H), 6.80-6.76 (m, 1 H), 2.18-
15 2.11 (m, 1 H), 1.90-1.81 (m, 1 H), 1.30-1.19 (m, 3 H), 0.97-0.80 (m, 4 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -82.692, -123.78-123.86; MS (CI) *m/e* calc'd for C₁₃H₁₅F₄N₂O: 291.112051, found 291.112227; 291 (MH⁺, 100).

20

Example 25

Preparation of (+/-)-6-Fluoro-4-phenylethyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Phenyl)

25 A solution of XVII-a (R⁴ = phenyl) (30 mg, 90 mmol) in ethanol (1 mL) and EtOAc (0.5 mL) was treated with 10% Pd on carbon under H₂ according to the procedure of Example 15 to afford 20 mg (67%) of the desired product. No further purification was necessary: mp 98 °C; ¹H NMR (300 MHz, acetone-d₆) δ 7.18-6.99 (m, 7H), 6.84-6.79 (m, 1 H), 2.68-2.60
30 (m, 1 H), 2.48-2.12 (m, 3 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -82.67, -123.24-123.32; MS (CI) *m/e* calc'd for C₁₇H₁₅F₄N₂O: 339.112051, found 339.110781; 339 (MH⁺, 100).

Example 26


Preparation of (+/-)-6-Fluoro-4-methylpropargyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (R⁴ = Methyl)

5

Synthesis of XIX-a from XVI-a.

A solution of 2-butyne (94 mg, 1.75 mmol) in anhydrous THF (3.5 mL) was cooled to 0 °C, treated with n-BuLi (0.97 mL of 1.6 M solution in hexanes, 1.55 mmol), and aged for 0.5 h. To a solution of XVI-a (90 mg, 0.388 mmol) in anhydrous THF (1.9 mL) at -78 °C was added the lithium anion over 5 minutes, followed by boron trifluoride etherate (25 mL, 0.194 mmol). The cooling bath was removed and the mixture was allowed to warm to room temperature. After 16 h at room temperature, quench by addition of 1 M citric acid (10 mL), dilute with EtOAc (50 mL), separate phases and wash the organic phase sequentially with saturated aqueous NaHCO₃ (20 mL) and saturated aqueous NaCl (20 mL). The resulting material was purified by HPLC (2.5% MeOH/CH₂Cl₂) to afford 10 mg (9%) of the desired product: mp 181 °C; ¹H NMR (300 MHz, acetone-d₆) δ 8.91 (br s, 1 H), 7.27 (d, 8.4H), 7.18-7.08 (m, 1 H), 7.02-6.97 (m, 2 H), 3.29 (dd, J = 16.8, 2.6 Hz, 1 H), 3.00 (dd, J = 16.8, 2.2 Hz, 1 H), 1.61-1.59 (m, 3 H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -81.86, -123.69-123.70; MS (CI) m/e calc'd for C₁₃H₁₁F₄N₂O: 287.080751, found 287.080340; 287 (MH⁺, 75), 304 (M+NH₄⁺, 100).

SCHEME 4: Chiral Resolution



5

Examples 27 and 28

Preparation of (-)-6-Chloro-4-cyclopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (Example 27) and (+)-6-Chloro-4-cyclopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (Example 28)

10

Resolution of IV-b,c from IV-a (R^4 = Cyclopropyl).

Chiral HPLC utilizing a Chiralcel OD column, 3% isopropanol, 5% CH_2Cl_2 and 92% hexanes at ambient temperature with a 1.0 mL/min flow rate and detection at 250 nm afforded separation of IV-b from IV-c with enantiomeric excesses of 99% and 99.4%, respectively. IV-b: mp 106-109 °C; $[\alpha]_{\text{D}}^{25}$ -60.34° (c=0.274, MeOH). IV-c: mp 105-107 °C; $[\alpha]_{\text{D}}^{25}$ +58.33° (c=0.288, MeOH).

20

Examples 29 and 30

Preparation of (+)-4-Cyclopropylethynyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (Example 29) and (-)-4-Cyclopropylethynyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (Example 30)

25

Resolution of XII-b,c from XII-a (R^4 = Cyclopropyl).

Chiral HPLC utilizing a Chiralpak AD column, 5% water and 95% methanol at ambient temperature with a 0.8 mL/min flow rate and detection at 250 nm afforded separation of XII-b from XII-c with enantiomeric excesses of 100% and 99%, respectively. XII-b: mp 187 °C; $[\alpha]_D^{25} +1.46^\circ$ (c=0.274, MeOH). XII-c: mp 187.5-188.8 °C; $[\alpha]_D^{25} -1.45^\circ$ (c=0.278, MeOH).

Examples 31 and 32

Preparation of (-)-5,6-Difluoro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (Example 31) and (+)-5,6-Difluoro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (Example 32)

Resolution of XII-b,c from XII-a (R^4 = Isopropyl).

Chiral HPLC utilizing a Chiralpak AD column, 5% water and 95% methanol at ambient temperature with a 0.5 mL/min flow rate and detection at 250 nm afforded separation of XII-b from XII-c with enantiomeric excesses of 100% and 99%, respectively. XII-b: mp 155 °C; $[\alpha]_D^{25} -2.14^\circ$ (c=0.280, MeOH). XII-c: 98 °C; $[\alpha]_D^{25} +4.45^\circ$ (c=0.292, MeOH).

Examples 33 and 34

Preparation of (-)-5,6-Difluoro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (Example 33) and (+)-5,6-Difluoro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (Example 34)

Resolution of XII-b,c from XII-a (R^4 = Ethyl).


Chiral HPLC utilizing a AS column, 20% ethanol and 80% hexanes at ambient temperature with a 1.0 mL/min flow rate and detection at 250 nm afforded separation of XII-b from

XII-c with enantiomeric excesses of 100% and 99%, respectively. XII-b: mp 165-167 °C. XII-c: mp 157-159 °C.

Examples 35 and 36

- 5 Preparation of 5,6-Difluoro-4-(2-hydroxyethyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (Example 35) and 5,6-Difluoro-4-(1-hydroxyethyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone (Example 36)

10



To a slurry of ketimine (300 mg, 1.20 mmol) in anhyd. THF (11 mL) at -78 °C was sequentially added a precooled (0 °C) solution of the silyl protected lithium acetylide (5.40 mmol) and $\text{BF}_3\cdot\text{OEt}_2$ (0.60 mmol). The resulting mixture was stirred at rt overnight. The reaction was quenched by the addition of 1 M citric acid and diluted with EtOAc. The phases were separated, the organic phase was washed with water, sat. aq. NaHCO_3 and sat. aq. NaCl . The organic extracts were dried over MgSO_4 , filtered and concentrated.

The material was purified by regular phase HPLC chromatography (41.4 mm Rainin Dynamax® column using 60 Å silica @ 25 mL/min): 2.5% MeOH/CH₂Cl₂ for 24 min, increase to 30% MeOH/CH₂Cl₂ over 4 min, 30% MeOH/CH₂Cl₂ for 10 min, and ramp back to 2.5% MeOH/CH₂Cl₂ over 2 min. The yield of the protected intermediates was 47% and 32%, respectively.

Example 35-intermediate. Mp 62.9-64 °C; ¹H NMR (300 MHz, acetone-d₆) δ 8.98 (br s, 1H), 7.41-7.32 (m, 2H), 6.83-6.78 (m, 1H), 3.74 (t, J = 6.6 Hz, 2H), 2.47 (t, J = 6.6 Hz, 2H), 0.81 (s, 9H), 0.00 (s, 6H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -83.17, -135.16--135.31, -148.09--148.22; MS (CI) calc'd for C₁₉H₂₄F₅N₂O₂Si: m/z 435.152723, found 435.151149; 435 (MH⁺, 94), 452 (M+NH₄⁺, 100); Analysis calc'd for C₁₉H₂₃F₅N₂O₂Si: C, 52.52; H, 5.35; N, 6.46; found: C, 52.65; H, 5.29; N, 6.31.

Example 36-intermediate. ¹H NMR (300 MHz, acetone-d₆) δ 8.96 (br s, 1H), 7.50 (br s, 1H), 7.37-7.28 (m, 1H), 6.79-6.74 (m, 1H), 4.61 (q, J = 13.2, 6.6 Hz, 1H), 1.30 (d, J = 6.6 Hz, 3H), 0.78 (s, 9H), 0.01 (s, 6H); ¹⁹F NMR (282 MHz, acetone-d₆) δ -82.88--82.95, -135.20--135.42, -148.06--148.23; MS (CI) calc'd for C₁₉H₂₄F₅N₂O₂Si: m/z 435.152723, found 435.152927; 435 (MH⁺, 51), 452 (M+NH₄⁺, 100); Analysis calc'd for C₁₉H₂₃F₅N₂O₂Si: C, 52.52; H, 5.35; N, 6.46; found: C, 52.54; H, 5.34; N, 6.69.

To a solution of the protected intermediate for Example 35 (0.56 mmol) in THF (1.1 mL) was added TBAF (0.62 mL of 1.0 M solution in THF). The resulting mixture was stirred at rt for 1 h, diluted with EtOAc, washed with 1 N HCl, sat. aq. NaHCO₃, and sat. aq. NaCl. The organic extract was dried over MgSO₄, filtered and concentrated. The material was purified by regular phase HPLC chromatography (41.4 mm Rainin Dynamax® column using 60 Å silica @ 25 mL/min): 2.5% MeOH/CH₂Cl₂ for 24 min, increase to 30% MeOH/CH₂Cl₂ over 4 min, 30% MeOH/CH₂Cl₂ for 10 min, and ramp back to 2.5% MeOH/CH₂Cl₂ over 2 min. Example 35 was isolated in 82% yield.

Example 35. Mp 190-192 °C; ¹H NMR (300 MHz, acetone-d₆) δ 9.05 (br s, 1H), 7.53 (br s, 1H), 7.45-7.36 (m, 1H), 6.88-6.83 (m, 1H), 4.01-3.98 (m, 1H), 3.68-3.64 (m, 2H), 2.50 (t,

J = 6.6 Hz, 2H); ^{19}F NMR (282 MHz, acetone- d_6) δ -83.3, -135.68--135.88, -148.10--148.22; MS (CI) calc'd for $\text{C}_{13}\text{H}_{10}\text{F}_5\text{N}_2\text{O}_2$: m/z 321.066244, found 321.066479; 321 (MH $^+$, 100); Analysis calc'd for $\text{C}_{13}\text{H}_9\text{F}_5\text{N}_2\text{O}_2$: C, 48.76; H, 2.83; N, 8.76; found: C, 49.05; H, 3.23; N, 8.38.

Example 36 was synthesized in an analogous manner to deliver the title compound in 88% yield. Mp 190-191 °C; ^1H NMR (300 MHz, acetone- d_6) δ 9.06 (br s, 1H), 7.56 (br s, 1H), 7.46-7.37 (m, 1H), 6.88-6.83 (m, 1H), 4.58-4.57 (m, 2H), 1.39 (d, J = 5.5 Hz, 3H); ^{19}F NMR (282 MHz, acetone- d_6) δ -83.15, -135.40, -135.60, -148.08--148.20; MS (CI) calc'd for $\text{C}_{13}\text{H}_{10}\text{F}_5\text{N}_2\text{O}_2$: m/z 321.066244, found 321.065983; 321 (MH $^+$, 58), 338 (M+NH $_4^+$, 100); Analysis calc'd for $\text{C}_{13}\text{H}_9\text{F}_5\text{N}_2\text{O}_2$: C, 48.76; H, 2.83; N, 8.76; found: C, 48.84; H, 2.76; N, 8.63.

Example 37

Preparation of (+)-4-E-Cyclopropylethenyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone

To a solution of XII-b (200 mg, 0.632 mmol) in anhyd. THF (1.3 mL) at rt was added a solution of lithium aluminium hydride (1.3 mL of 1.0 M solution in THF). The resulting mixture was stirred at rt overnight. The reaction was quenched by addition of 10% NaOH (3 mL) and water (3 mL). The mixture was diluted with EtOAc (30 mL) and the phases were separated. The organic phase was washed with sat. aq. NaCl, dried over MgSO_4 , filtered and concentrated. The title compound was purified by regular phase HPLC (41.4 mm Rainin Dynamax® column using 60 Å silica): 2.5% MeOH/ CH_2Cl_2 for 24 min, increase to 30% MeOH/ CH_2Cl_2 over 4 min, 30% MeOH/ CH_2Cl_2 for 10 min, and ramp back to 2.5% MeOH/ CH_2Cl_2 over 2 min. Mp 80-83 °C; ^1H NMR (300 MHz, acetone- d_6) δ 9.07 (br s, 1H), 7.33 (q, J=8.8 Hz, 1H), 6.94 (br s, 1H), 6.84-6.79 (m, 1H), 6.27 (dd, J = 15.6, 7.5 Hz, 1H), 5.67 (dd, J = 15.2, 9.4 Hz, 1H), 1.65-1.56 (m, 1H), 0.80-0.71 (m, 2H), 0.50-0.42 (m, 2H); ^{19}F NMR (282 MHz, acetone- d_6) δ -82.68, -135.05, -148.49; MS (CI) calc'd for $\text{C}_{14}\text{H}_{12}\text{F}_5\text{N}_2\text{O}$: m/z 319.086979, found


319.087755; 319 (MH⁺, 100); [α]_D²⁰ +72.77° (c=0.382, MeOH);
Analysis calc'd for C₁₄H₁₁F₅N₂O: C, 52.84; H, 3.48; N,
8.80; found: C, 53.02; H, 3.48; N, 8.61.

5

Example 38**Preparation of (-)-6-Chloro-4-E-cyclopropylethenyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone**

The title compound was prepared as described for Example
10 37 (starting from IV-b), except that it was purified using a
Chiralcel OD column at 1.5 mL/min in 0.5% EtOH/20%
CH₂Cl₂/79.5% hexanes. Mp 87-89 °C; ¹H NMR (300 MHz, acetone-
d₆) δ 9.08 (br s, 1H), 7.40-7.25 (m, 2H), 7.04-6.90 (m, 2H),
6.28-6.18 (m, 1H), 5.64-5.52 (m, 1H), 1.68-1.55 (m, 1H),
15 0.83-0.71 (m, 2H), 0.53-0.41 (m, 2H); ¹⁹F NMR (282 MHz,
acetone-d₆) δ -81.67; MS (CI) calc'd for C₁₄H₁₃ClF₃N₂O: m/z
317.066851, found 317.065857; 317 (MH⁺, 100); [α]_D²⁰ -6.81°
(c=0.382, MeOH); Analysis calc'd for C₁₄H₁₂ClF₃N₂O . 0.27
C₃H₆O: C, 53.52; H, 4.13; N, 8.43; found: C, 53.90; H,
20 4.07; N, 8.80.

Table 1*




Ex. #	R ³	R ¹	R ²	R ⁸	m.p. (°C)	Mass Spec
1	6-Cl	CF ₃	C≡C-cycPr	H	86.6-88	332 (M+NH ₄ ⁺)
2	6-Cl	CF ₃	C≡C-iPr	H	180	334 (M+NH ₄ ⁺)
3	6-Cl	CF ₃	C≡C-2-Pyridyl	H	105	352 (MH ⁺)
4	6-Cl	CF ₃	C≡C-Et	H	217-219	303 (MH ⁺)
5	6-Cl	CF ₃	C≡C-Ph	H	104-107	368 (M+NH ₄ ⁺)
6	6-MeO	CF ₃	C≡C-cycPr	H	208	311 (MH ⁺)
7	6-MeO	CF ₃	C≡C-iPr	H	228-229	313 (MH ⁺)
8	6-MeO	CF ₃	C≡C-2-Pyridyl	H	97-98	348 (MH ⁺)
9	6-MeO	CF ₃	C≡C-Ph	H	206.2-207.7	347 (MH ⁺)
10	5,6-diF	CF ₃	C≡C-cycPr	H	101 dec.	317 (MH ⁺)
11	5,6-diF	CF ₃	C≡C-iPr	H	79-80	319 (MH ⁺)
12	5,6-diF	CF ₃	C≡C-2-Pyridyl	H	219-220	354 (MH ⁺)
13	5,6-diF	CF ₃	C≡C-Et	H	191-194	305 (MH ⁺)
14	5,6-diF	CF ₃	C≡C-Ph	H	215-217	370 (M+NH ₄ ⁺)
15	5,6-diF	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	H	192-193	323 (MH ⁺)
16	5,6-diF	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	H		309 (MH ⁺)
17	6-F	CF ₃	C≡C-cycPr	H	155	299 (MH ⁺)
18	6-F	CF ₃	C≡C-iPr	H	158	301 (MH ⁺)
19	6-F	CF ₃	C≡C-2-Pyridyl	H	155	336 (MH ⁺)
20	6-F	CF ₃	C≡C-Et	H	190	287 (MH ⁺)
21	6-F	CF ₃	C≡C-Ph	H	107	352 (M+NH ₄ ⁺)
22	6-F	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	H	179	305 (MH ⁺)

23	6-F	CF ₃	CH ₂ CH ₂ -2-Pyridyl	H	88	340 (MH ⁺)
24	6-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	H	198	291 (MH ⁺)
25	6-F	CF ₃	CH ₂ CH ₂ Ph	H	98	339 (MH ⁺)
26	6-F	CF ₃	CH ₂ C≡C-CH ₃	H	181	304 (M+NH ₄ ⁺)
27(-)	6-Cl	CF ₃	C≡C-cycPr	H	106- 109	313 (M ⁻)
28(+)	6-Cl	CF ₃	C≡C-cycPr	H	105- 107	313 (M ⁻)
29(+)	5,6-diF	CF ₃	C≡C-cycPr	H	187	315 (M ⁻)
30(-)	5,6-diF	CF ₃	C≡C-cycPr	H	188- 189	315 (M ⁻)
31(-)	5,6-diF	CF ₃	C≡C-iPr	H	155	317 (M ⁻)
32(+)	5,6-diF	CF ₃	C≡C-iPr	H	98	317 (M ⁻)
33(-)	5,6-diF	CF ₃	C≡C-Et	H	165- 167	303 (M ⁻)
34(+)	5,6-diF	CF ₃	C≡C-Et	H	157- 159	303 (M ⁻)
35	5,6-diF	CF ₃	C≡CCH ₂ CH ₂ OH	H	190- 192	321 (MH ⁺)
36	5,6-diF	CF ₃	C≡C-CH(OH)Me	H	190- 191	338 (M+NH ₄ ⁺)
37(+)	5,6-diF	CF ₃	C=C-cycPr (E)	H	80-83	319 (MH ⁺)
38(-)	6-Cl	CF ₃	C=C-cycPr (E)	H	87-89	317 (MH ⁺)

*Unless otherwise indicated, stereochemistry is (+/-).

Table 2*



Ex. #	R ³	R ¹	R ²	R ⁸
1	6-Cl	CF ₃	C≡CCH ₂ CH ₂ OH	H
2	6-Cl	CF ₃	C≡C-CH(OH)Me	H
3	6-Cl	CF ₃	C≡C-(2-Cl)Ph	H
4	6-Cl	CF ₃	C≡C-(3-Cl)Ph	H
5	6-Cl	CF ₃	C≡C-(4-Cl)Ph	H
6	6-Cl	CF ₃	C≡C-(2-F)Ph	H
7	6-Cl	CF ₃	C≡C-(3-F)Ph	H
8	6-Cl	CF ₃	C≡C-(4-F)Ph	H
9	6-Cl	CF ₃	C≡C-(2-OH)Ph	H
10	6-Cl	CF ₃	C≡C-(3-OH)Ph	H
11	6-Cl	CF ₃	C≡C-(4-OH)Ph	H
12	6-Cl	CF ₃	C≡C-(2-OMe)Ph	H
13	6-Cl	CF ₃	C≡C-(3-OMe)Ph	H
14	6-Cl	CF ₃	C≡C-(4-OMe)Ph	H
15	6-Cl	CF ₃	C≡C-(2-CN)Ph	H
16	6-Cl	CF ₃	C≡C-(3-CN)Ph	H
17	6-Cl	CF ₃	C≡C-(4-CN)Ph	H
18	6-Cl	CF ₃	C≡C-(2-NO ₂)Ph	H
19	6-Cl	CF ₃	C≡C-(3-NO ₂)Ph	H
20	6-Cl	CF ₃	C≡C-(4-NO ₂)Ph	H
21	6-Cl	CF ₃	C≡C-(2-NH ₂)Ph	H
22	6-Cl	CF ₃	C≡C-(3-NH ₂)Ph	H
23	6-Cl	CF ₃	C≡C-(4-NH ₂)Ph	H
24	6-Cl	CF ₃	C≡C-(2-NMe ₂)Ph	H
25	6-Cl	CF ₃	C≡C-(3-NMe ₂)Ph	H
26	6-Cl	CF ₃	C≡C-(4-NMe ₂)Ph	H
27	6-Cl	CF ₃	C≡C-3-Pyridyl	H
28	6-Cl	CF ₃	C≡C-4-Pyridyl	H

29	6-Cl	CF ₃	C≡C-2-furanyl	H
30	6-Cl	CF ₃	C≡C-3-furanyl	H
31	6-Cl	CF ₃	C≡C-2-thienyl	H
32	6-Cl	CF ₃	C≡C-3-thienyl	H
33	6-Cl	CF ₃	C≡C-2-oxazolyl	H
34	6-Cl	CF ₃	C≡C-2-thiazolyl	H
35	6-Cl	CF ₃	C≡C-4-isoxazolyl	H
36	6-Cl	CF ₃	C≡C-2-imidazolyl	H
37	6-Cl	CF ₃	C=CCH ₂ CH ₂ OH	H
38	6-Cl	CF ₃	C=C-CH(OH)Me	H
39	6-Cl	CF ₃	C=C-(2-Cl)Ph	H
40	6-Cl	CF ₃	C=C-(3-Cl)Ph	H
41	6-Cl	CF ₃	C=C-(4-Cl)Ph	H
42	6-Cl	CF ₃	C=C-(2-F)Ph	H
43	6-Cl	CF ₃	C=C-(3-F)Ph	H
44	6-Cl	CF ₃	C=C-(4-F)Ph	H
45	6-Cl	CF ₃	C=C-(2-OH)Ph	H
46	6-Cl	CF ₃	C=C-(3-OH)Ph	H
47	6-Cl	CF ₃	C=C-(4-OH)Ph	H
48	6-Cl	CF ₃	C=C-(2-OMe)Ph	H
49	6-Cl	CF ₃	C=C-(3-OMe)Ph	H
50	6-Cl	CF ₃	C=C-(4-OMe)Ph	H
51	6-Cl	CF ₃	C=C-(2-CN)Ph	H
52	6-Cl	CF ₃	C=C-(3-CN)Ph	H
53	6-Cl	CF ₃	C=C-(4-CN)Ph	H
54	6-Cl	CF ₃	C=C-(2-NO ₂)Ph	H
55	6-Cl	CF ₃	C=C-(3-NO ₂)Ph	H
56	6-Cl	CF ₃	C=C-(4-NO ₂)Ph	H
57	6-Cl	CF ₃	C=C-(2-NH ₂)Ph	H
58	6-Cl	CF ₃	C=C-(3-NH ₂)Ph	H
59	6-Cl	CF ₃	C=C-(4-NH ₂)Ph	H
60	6-Cl	CF ₃	C=C-(2-NMe ₂)Ph	H
61	6-Cl	CF ₃	C=C-(3-NMe ₂)Ph	H
62	6-Cl	CF ₃	C=C-(4-NMe ₂)Ph	H

63	6-Cl	CF ₃	C=C-3-Pyridyl	H
64	6-Cl	CF ₃	C=C-4-Pyridyl	H
65	6-Cl	CF ₃	C=C-2-furanyl	H
66	6-Cl	CF ₃	C=C-3-furanyl	H
67	6-Cl	CF ₃	C=C-2-thienyl	H
68	6-Cl	CF ₃	C=C-3-thienyl	H
69	6-Cl	CF ₃	C=C-2-oxazolyl	H
70	6-Cl	CF ₃	C=C-2-thiazolyl	H
71	6-Cl	CF ₃	C=C-4-isoxazolyl	H
72	6-Cl	CF ₃	C=C-2-imidazolyl	H
73	6-Cl	CF ₃	CH ₂ CH ₂ -cycPr	H
74	6-Cl	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ OH	H
75	6-Cl	CF ₃	CH ₂ CH ₂ -CH(OH)Me	H
76	6-Cl	CF ₃	CH ₂ CH ₂ -Ph	H
77	6-Cl	CF ₃	CH ₂ CH ₂ -(2-Cl) Ph	H
78	6-Cl	CF ₃	CH ₂ CH ₂ -(3-Cl) Ph	H
79	6-Cl	CF ₃	CH ₂ CH ₂ -(4-Cl) Ph	H
80	6-Cl	CF ₃	CH ₂ CH ₂ -(2-F) Ph	H
81	6-Cl	CF ₃	CH ₂ CH ₂ -(3-F) Ph	H
82	6-Cl	CF ₃	CH ₂ CH ₂ -(4-F) Ph	H
83	6-Cl	CF ₃	CH ₂ CH ₂ -(2-OH) Ph	H
84	6-Cl	CF ₃	CH ₂ CH ₂ -(3-OH) Ph	H
85	6-Cl	CF ₃	CH ₂ CH ₂ -(4-OH) Ph	H
86	6-Cl	CF ₃	CH ₂ CH ₂ -(2-OMe) Ph	H
87	6-Cl	CF ₃	CH ₂ CH ₂ -(3-OMe) Ph	H
88	6-Cl	CF ₃	CH ₂ CH ₂ -(4-OMe) Ph	H
89	6-Cl	CF ₃	CH ₂ CH ₂ -(2-CN) Ph	H
90	6-Cl	CF ₃	CH ₂ CH ₂ -(3-CN) Ph	H
91	6-Cl	CF ₃	CH ₂ CH ₂ -(4-CN) Ph	H
92	6-Cl	CF ₃	CH ₂ CH ₂ -(2-NO ₂) Ph	H
93	6-Cl	CF ₃	CH ₂ CH ₂ -(3-NO ₂) Ph	H
94	6-Cl	CF ₃	CH ₂ CH ₂ -(4-NO ₂) Ph	H
95	6-Cl	CF ₃	CH ₂ CH ₂ -(2-NH ₂) Ph	H
96	6-Cl	CF ₃	CH ₂ CH ₂ -(3-NH ₂) Ph	H

97	6-Cl	CF ₃	CH ₂ CH ₂ -(4-NH ₂) Ph	H
98	6-Cl	CF ₃	CH ₂ CH ₂ -(2-NMe ₂) Ph	H
99	6-Cl	CF ₃	CH ₂ CH ₂ -(3-NMe ₂) Ph	H
100	6-Cl	CF ₃	CH ₂ CH ₂ -(4-NMe ₂) Ph	H
101	6-Cl	CF ₃	CH ₂ CH ₂ -2-Pyridyl	H
102	6-Cl	CF ₃	CH ₂ CH ₂ -3-Pyridyl	H
103	6-Cl	CF ₃	CH ₂ CH ₂ -4-Pyridyl	H
104	6-Cl	CF ₃	CH ₂ CH ₂ -2-furanyl	H
105	6-Cl	CF ₃	CH ₂ CH ₂ -3-furanyl	H
106	6-Cl	CF ₃	CH ₂ CH ₂ -4-furanyl	H
107	6-Cl	CF ₃	CH ₂ CH ₂ -3-thienyl	H
108	6-Cl	CF ₃	CH ₂ CH ₂ -2-oxazolyl	H
109	6-Cl	CF ₃	CH ₂ CH ₂ -2-thiazolyl	H
110	6-Cl	CF ₃	CH ₂ CH ₂ -4-isoxazolyl	H
111	6-Cl	CF ₃	CH ₂ CH ₂ -2-imidazolyl	H
112	6-Cl	CF ₃	C≡C-cycPr	CH ₃
113	6-Cl	CF ₃	C≡C-Ph	CH ₃
114	6-Cl	CF ₃	C≡C-2-Pyridyl	CH ₃
115	6-Cl	CF ₃	C≡C-3-Pyridyl	CH ₃
116	6-Cl	CF ₃	C≡C-4-Pyridyl	CH ₃
117	6-Cl	CF ₃	C≡C-2-furanyl	CH ₃
118	6-Cl	CF ₃	C≡C-3-furanyl	CH ₃
119	6-Cl	CF ₃	C≡C-2-thienyl	CH ₃
120	6-Cl	CF ₃	C≡C-3-thienyl	CH ₃
121	6-Cl	CF ₃	C=C-cycPr	CH ₃
122	6-Cl	CF ₃	C=C-Ph	CH ₃
123	6-Cl	CF ₃	C=C-2-Pyridyl	CH ₃
124	6-Cl	CF ₃	C=C-3-Pyridyl	CH ₃
125	6-Cl	CF ₃	C=C-4-Pyridyl	CH ₃
126	6-Cl	CF ₃	C=C-2-furanyl	CH ₃
127	6-Cl	CF ₃	C=C-3-furanyl	CH ₃
128	6-Cl	CF ₃	C=C-2-thienyl	CH ₃
129	6-Cl	CF ₃	C=C-3-thienyl	CH ₃
130	6-Cl	CF ₃	CH ₂ CH ₂ -cycPr	CH ₃

131	6-Cl	CF ₃	CH ₂ CH ₂ -Ph	CH ₃
132	6-Cl	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₃
133	6-Cl	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₃
134	6-Cl	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₃
135	6-Cl	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₃
136	6-Cl	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₃
137	6-Cl	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₃
138	6-Cl	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₃
139	6-Cl	CF ₃	C≡C-cycPr	CH ₂ CH ₃
140	6-Cl	CF ₃	C≡C-Ph	CH ₂ CH ₃
141	6-Cl	CF ₃	C≡C-2-Pyridyl	CH ₂ CH ₃
142	6-Cl	CF ₃	C≡C-3-Pyridyl	CH ₂ CH ₃
143	6-Cl	CF ₃	C≡C-4-Pyridyl	CH ₂ CH ₃
144	6-Cl	CF ₃	C≡C-2-furanyl	CH ₂ CH ₃
145	6-Cl	CF ₃	C≡C-3-furanyl	CH ₂ CH ₃
146	6-Cl	CF ₃	C≡C-2-thienyl	CH ₂ CH ₃
147	6-Cl	CF ₃	C≡C-3-thienyl	CH ₂ CH ₃
148	6-Cl	CF ₃	C=C-cycPr	CH ₂ CH ₃
149	6-Cl	CF ₃	C=C-Ph	CH ₂ CH ₃
150	6-Cl	CF ₃	C=C-2-Pyridyl	CH ₂ CH ₃
151	6-Cl	CF ₃	C=C-3-Pyridyl	CH ₂ CH ₃
152	6-Cl	CF ₃	C=C-4-Pyridyl	CH ₂ CH ₃
153	6-Cl	CF ₃	C=C-2-furanyl	CH ₂ CH ₃
154	6-Cl	CF ₃	C=C-3-furanyl	CH ₂ CH ₃
155	6-Cl	CF ₃	C=C-2-thienyl	CH ₂ CH ₃
156	6-Cl	CF ₃	C=C-3-thienyl	CH ₂ CH ₃
157	6-Cl	CF ₃	CH ₂ CH ₂ -cycPr	CH ₂ CH ₃
158	6-Cl	CF ₃	CH ₂ CH ₂ -Ph	CH ₂ CH ₃
159	6-Cl	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₂ CH ₃
160	6-Cl	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₂ CH ₃
161	6-Cl	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₂ CH ₃
162	6-Cl	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₂ CH ₃
163	6-Cl	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₂ CH ₃
164	6-Cl	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₂ CH ₃

165	6-Cl	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₂ CH ₃
166	6-MeO	CF ₃	C≡CCH ₂ CH ₂ OH	H
167	6-MeO	CF ₃	C≡C-CH(OH)Me	H
168	6-MeO	CF ₃	C≡C-(2-Cl)Ph	H
169	6-MeO	CF ₃	C≡C-(3-Cl)Ph	H
170	6-MeO	CF ₃	C≡C-(4-Cl)Ph	H
171	6-MeO	CF ₃	C≡C-(2-F)Ph	H
172	6-MeO	CF ₃	C≡C-(3-F)Ph	H
173	6-MeO	CF ₃	C≡C-(4-F)Ph	H
174	6-MeO	CF ₃	C≡C-(2-OH)Ph	H
175	6-MeO	CF ₃	C≡C-(3-OH)Ph	H
176	6-MeO	CF ₃	C≡C-(4-OH)Ph	H
177	6-MeO	CF ₃	C≡C-(2-OMe)Ph	H
178	6-MeO	CF ₃	C≡C-(3-OMe)Ph	H
179	6-MeO	CF ₃	C≡C-(4-OMe)Ph	H
180	6-MeO	CF ₃	C≡C-(2-CN)Ph	H
181	6-MeO	CF ₃	C≡C-(3-CN)Ph	H
182	6-MeO	CF ₃	C≡C-(4-CN)Ph	H
183	6-MeO	CF ₃	C≡C-(2-NO ₂)Ph	H
184	6-MeO	CF ₃	C≡C-(3-NO ₂)Ph	H
185	6-MeO	CF ₃	C≡C-(4-NO ₂)Ph	H
186	6-MeO	CF ₃	C≡C-(2-NH ₂)Ph	H
187	6-MeO	CF ₃	C≡C-(3-NH ₂)Ph	H
188	6-MeO	CF ₃	C≡C-(4-NH ₂)Ph	H
189	6-MeO	CF ₃	C≡C-(2-NMe ₂)Ph	H
190	6-MeO	CF ₃	C≡C-(3-NMe ₂)Ph	H
191	6-MeO	CF ₃	C≡C-(4-NMe ₂)Ph	H
192	6-MeO	CF ₃	C≡C-3-Pyridyl	H
193	6-MeO	CF ₃	C≡C-4-Pyridyl	H
194	6-MeO	CF ₃	C≡C-2-furanyl	H
195	6-MeO	CF ₃	C≡C-3-furanyl	H
196	6-MeO	CF ₃	C≡C-2-thienyl	H
197	6-MeO	CF ₃	C≡C-3-thienyl	H
198	6-MeO	CF ₃	C≡C-2-oxazolyl	H

199	6-MeO	CF ₃	C≡C-2-thiazolyl	H
200	6-MeO	CF ₃	C≡C-4-isoxazolyl	H
201	6-MeO	CF ₃	C≡C-2-imidazolyl	H
202	6-MeO	CF ₃	C=CCH ₂ CH ₂ OH	H
203	6-MeO	CF ₃	C=C-CH(OH)Me	H
204	6-MeO	CF ₃	C=C-(2-Cl)Ph	H
205	6-MeO	CF ₃	C=C-(3-Cl)Ph	H
206	6-MeO	CF ₃	C=C-(4-Cl)Ph	H
207	6-MeO	CF ₃	C=C-(2-F)Ph	H
208	6-MeO	CF ₃	C=C-(3-F)Ph	H
209	6-MeO	CF ₃	C=C-(4-F)Ph	H
210	6-MeO	CF ₃	C=C-(2-OH)Ph	H
211	6-MeO	CF ₃	C=C-(3-OH)Ph	H
212	6-MeO	CF ₃	C=C-(4-OH)Ph	H
213	6-MeO	CF ₃	C=C-(2-OMe)Ph	H
214	6-MeO	CF ₃	C=C-(3-OMe)Ph	H
215	6-MeO	CF ₃	C=C-(4-OMe)Ph	H
216	6-MeO	CF ₃	C=C-(2-CN)Ph	H
217	6-MeO	CF ₃	C=C-(3-CN)Ph	H
218	6-MeO	CF ₃	C=C-(4-CN)Ph	H
219	6-MeO	CF ₃	C=C-(2-NO ₂)Ph	H
220	6-MeO	CF ₃	C=C-(3-NO ₂)Ph	H
221	6-MeO	CF ₃	C=C-(4-NO ₂)Ph	H
222	6-MeO	CF ₃	C=C-(2-NH ₂)Ph	H
223	6-MeO	CF ₃	C=C-(3-NH ₂)Ph	H
224	6-MeO	CF ₃	C=C-(4-NH ₂)Ph	H
225	6-MeO	CF ₃	C=C-(2-NMe ₂)Ph	H
226	6-MeO	CF ₃	C=C-(3-NMe ₂)Ph	H
227	6-MeO	CF ₃	C=C-(4-NMe ₂)Ph	H
228	6-MeO	CF ₃	C=C-3-Pyridyl	H
229	6-MeO	CF ₃	C=C-4-Pyridyl	H
230	6-MeO	CF ₃	C=C-2-furanyl	H
231	6-MeO	CF ₃	C=C-3-furanyl	H
232	6-MeO	CF ₃	C=C-2-thienyl	H

233	6-MeO	CF ₃	C=C-3-thienyl	H
234	6-MeO	CF ₃	C=C-2-oxazolyl	H
235	6-MeO	CF ₃	C=C-2-thiazolyl	H
236	6-MeO	CF ₃	C=C-4-isoxazolyl	H
237	6-MeO	CF ₃	C=C-2-imidazolyl	H
238	6-MeO	CF ₃	CH ₂ CH ₂ -cycPr	H
239	6-MeO	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ OH	H
240	6-MeO	CF ₃	CH ₂ CH ₂ -CH(OH)Me	H
241	6-MeO	CF ₃	CH ₂ CH ₂ -Ph	H
242	6-MeO	CF ₃	CH ₂ CH ₂ -(2-Cl)Ph	H
243	6-MeO	CF ₃	CH ₂ CH ₂ -(3-Cl)Ph	H
244	6-MeO	CF ₃	CH ₂ CH ₂ -(4-Cl)Ph	H
245	6-MeO	CF ₃	CH ₂ CH ₂ -(2-F)Ph	H
246	6-MeO	CF ₃	CH ₂ CH ₂ -(3-F)Ph	H
247	6-MeO	CF ₃	CH ₂ CH ₂ -(4-F)Ph	H
248	6-MeO	CF ₃	CH ₂ CH ₂ -(2-OH)Ph	H
249	6-MeO	CF ₃	CH ₂ CH ₂ -(3-OH)Ph	H
250	6-MeO	CF ₃	CH ₂ CH ₂ -(4-OH)Ph	H
251	6-MeO	CF ₃	CH ₂ CH ₂ -(2-OMe)Ph	H
252	6-MeO	CF ₃	CH ₂ CH ₂ -(3-OMe)Ph	H
253	6-MeO	CF ₃	CH ₂ CH ₂ -(4-OMe)Ph	H
254	6-MeO	CF ₃	CH ₂ CH ₂ -(2-CN)Ph	H
255	6-MeO	CF ₃	CH ₂ CH ₂ -(3-CN)Ph	H
256	6-MeO	CF ₃	CH ₂ CH ₂ -(4-CN)Ph	H
257	6-MeO	CF ₃	CH ₂ CH ₂ -(2-NO ₂)Ph	H
258	6-MeO	CF ₃	CH ₂ CH ₂ -(3-NO ₂)Ph	H
259	6-MeO	CF ₃	CH ₂ CH ₂ -(4-NO ₂)Ph	H
260	6-MeO	CF ₃	CH ₂ CH ₂ -(2-NH ₂)Ph	H
261	6-MeO	CF ₃	CH ₂ CH ₂ -(3-NH ₂)Ph	H
262	6-MeO	CF ₃	CH ₂ CH ₂ -(4-NH ₂)Ph	H
263	6-MeO	CF ₃	CH ₂ CH ₂ -(2-NMe ₂)Ph	H
264	6-MeO	CF ₃	CH ₂ CH ₂ -(3-NMe ₂)Ph	H
265	6-MeO	CF ₃	CH ₂ CH ₂ -(4-NMe ₂)Ph	H
266	6-MeO	CF ₃	CH ₂ CH ₂ -2-Pyridyl	H

267	6-MeO	CF ₃	CH ₂ CH ₂ -3-Pyridyl	H
268	6-MeO	CF ₃	CH ₂ CH ₂ -4-Pyridyl	H
269	6-MeO	CF ₃	CH ₂ CH ₂ -2-furanyl	H
270	6-MeO	CF ₃	CH ₂ CH ₂ -3-furanyl	H
271	6-MeO	CF ₃	CH ₂ CH ₂ -4-furanyl	H
272	6-MeO	CF ₃	CH ₂ CH ₂ -3-thienyl	H
273	6-MeO	CF ₃	CH ₂ CH ₂ -2-oxazolyl	H
274	6-MeO	CF ₃	CH ₂ CH ₂ -2-thiazolyl	H
275	6-MeO	CF ₃	CH ₂ CH ₂ -4-isoxazolyl	H
276	6-MeO	CF ₃	CH ₂ CH ₂ -2-imidazolyl	H
277	6-MeO	CF ₃	C≡C-cycPr	CH ₃
278	6-MeO	CF ₃	C≡C-Ph	CH ₃
279	6-MeO	CF ₃	C≡C-2-Pyridyl	CH ₃
280	6-MeO	CF ₃	C≡C-3-Pyridyl	CH ₃
281	6-MeO	CF ₃	C≡C-4-Pyridyl	CH ₃
282	6-MeO	CF ₃	C≡C-2-furanyl	CH ₃
283	6-MeO	CF ₃	C≡C-3-furanyl	CH ₃
284	6-MeO	CF ₃	C≡C-2-thienyl	CH ₃
285	6-MeO	CF ₃	C≡C-3-thienyl	CH ₃
286	6-MeO	CF ₃	C=C-cycPr	CH ₃
287	6-MeO	CF ₃	C=C-Ph	CH ₃
288	6-MeO	CF ₃	C=C-2-Pyridyl	CH ₃
289	6-MeO	CF ₃	C=C-3-Pyridyl	CH ₃
290	6-MeO	CF ₃	C=C-4-Pyridyl	CH ₃
291	6-MeO	CF ₃	C=C-2-furanyl	CH ₃
292	6-MeO	CF ₃	C=C-3-furanyl	CH ₃
293	6-MeO	CF ₃	C=C-2-thienyl	CH ₃
294	6-MeO	CF ₃	C=C-3-thienyl	CH ₃
295	6-MeO	CF ₃	CH ₂ CH ₂ -cycPr	CH ₃
296	6-MeO	CF ₃	CH ₂ CH ₂ -Ph	CH ₃
297	6-MeO	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₃
298	6-MeO	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₃
299	6-MeO	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₃
300	6-MeO	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₃

301	6-MeO	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₃
302	6-MeO	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₃
303	6-MeO	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₃
304	6-MeO	CF ₃	C≡C-cycPr	CH ₂ CH ₃
305	6-MeO	CF ₃	C≡C-Ph	CH ₂ CH ₃
306	6-MeO	CF ₃	C≡C-2-Pyridyl	CH ₂ CH ₃
307	6-MeO	CF ₃	C≡C-3-Pyridyl	CH ₂ CH ₃
308	6-MeO	CF ₃	C≡C-4-Pyridyl	CH ₂ CH ₃
309	6-MeO	CF ₃	C≡C-2-furanyl	CH ₂ CH ₃
310	6-MeO	CF ₃	C≡C-3-furanyl	CH ₂ CH ₃
311	6-MeO	CF ₃	C≡C-2-thienyl	CH ₂ CH ₃
312	6-MeO	CF ₃	C≡C-3-thienyl	CH ₂ CH ₃
313	6-MeO	CF ₃	C=C-cycPr	CH ₂ CH ₃
314	6-MeO	CF ₃	C=C-Ph	CH ₂ CH ₃
315	6-MeO	CF ₃	C=C-2-Pyridyl	CH ₂ CH ₃
316	6-MeO	CF ₃	C=C-3-Pyridyl	CH ₂ CH ₃
317	6-MeO	CF ₃	C=C-4-Pyridyl	CH ₂ CH ₃
318	6-MeO	CF ₃	C=C-2-furanyl	CH ₂ CH ₃
319	6-MeO	CF ₃	C=C-3-furanyl	CH ₂ CH ₃
320	6-MeO	CF ₃	C=C-2-thienyl	CH ₂ CH ₃
321	6-MeO	CF ₃	C=C-3-thienyl	CH ₂ CH ₃
322	6-MeO	CF ₃	CH ₂ CH ₂ -cycPr	CH ₂ CH ₃
323	6-MeO	CF ₃	CH ₂ CH ₂ -Ph	CH ₂ CH ₃
324	6-MeO	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₂ CH ₃
325	6-MeO	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₂ CH ₃
326	6-MeO	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₂ CH ₃
327	6-MeO	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₂ CH ₃
328	6-MeO	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₂ CH ₃
329	6-MeO	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₂ CH ₃
330	6-MeO	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₂ CH ₃
331	5,6-diF	CF ₃	C≡C-(2-Cl) Ph	H
332	5,6-diF	CF ₃	C≡C-(3-Cl) Ph	H
333	5,6-diF	CF ₃	C≡C-(4-Cl) Ph	H
334	5,6-diF	CF ₃	C≡C-(2-F) Ph	H

335	5,6-diF	CF ₃	C≡C-(3-F) Ph	H
336	5,6-diF	CF ₃	C≡C-(4-F) Ph	H
337	5,6-diF	CF ₃	C≡C-(2-OH) Ph	H
338	5,6-diF	CF ₃	C≡C-(3-OH) Ph	H
339	5,6-diF	CF ₃	C≡C-(4-OH) Ph	H
340	5,6-diF	CF ₃	C≡C-(2-OMe) Ph	H
341	5,6-diF	CF ₃	C≡C-(3-OMe) Ph	H
342	5,6-diF	CF ₃	C≡C-(4-OMe) Ph	H
343	5,6-diF	CF ₃	C≡C-(2-CN) Ph	H
344	5,6-diF	CF ₃	C≡C-(3-CN) Ph	H
345	5,6-diF	CF ₃	C≡C-(4-CN) Ph	H
346	5,6-diF	CF ₃	C≡C-(2-NO ₂) Ph	H
347	5,6-diF	CF ₃	C≡C-(3-NO ₂) Ph	H
348	5,6-diF	CF ₃	C≡C-(4-NO ₂) Ph	H
349	5,6-diF	CF ₃	C≡C-(2-NH ₂) Ph	H
350	5,6-diF	CF ₃	C≡C-(3-NH ₂) Ph	H
351	5,6-diF	CF ₃	C≡C-(4-NH ₂) Ph	H
352	5,6-diF	CF ₃	C≡C-(2-NMe ₂) Ph	H
353	5,6-diF	CF ₃	C≡C-(3-NMe ₂) Ph	H
354	5,6-diF	CF ₃	C≡C-(4-NMe ₂) Ph	H
355	5,6-diF	CF ₃	C≡C-3-Pyridyl	H
356	5,6-diF	CF ₃	C≡C-4-Pyridyl	H
357	5,6-diF	CF ₃	C≡C-2-furanyl	H
358	5,6-diF	CF ₃	C≡C-3-furanyl	H
359	5,6-diF	CF ₃	C≡C-2-thienyl	H
360	5,6-diF	CF ₃	C≡C-3-thienyl	H
361	5,6-diF	CF ₃	C≡C-2-oxazolyl	H
362	5,6-diF	CF ₃	C≡C-2-thiazolyl	H
363	5,6-diF	CF ₃	C≡C-4-isoxazolyl	H
364	5,6-diF	CF ₃	C≡C-2-imidazolyl	H
365	5,6-diF	CF ₃	C=C-(2-Cl) Ph	H
366	5,6-diF	CF ₃	C=C-(3-Cl) Ph	H
367	5,6-diF	CF ₃	C=C-(4-Cl) Ph	H
368	5,6-diF	CF ₃	C=C-(2-F) Ph	H

369	5,6-diF	CF ₃	C=C-(3-F) Ph	H
370	5,6-diF	CF ₃	C=C-(4-F) Ph	H
371	5,6-diF	CF ₃	C=C-(2-OH) Ph	H
372	5,6-diF	CF ₃	C=C-(3-OH) Ph	H
373	5,6-diF	CF ₃	C=C-(4-OH) Ph	H
374	5,6-diF	CF ₃	C=C-(2-OMe) Ph	H
375	5,6-diF	CF ₃	C=C-(3-OMe) Ph	H
376	5,6-diF	CF ₃	C=C-(4-OMe) Ph	H
377	5,6-diF	CF ₃	C=C-(2-CN) Ph	H
378	5,6-diF	CF ₃	C=C-(3-CN) Ph	H
379	5,6-diF	CF ₃	C=C-(4-CN) Ph	H
380	5,6-diF	CF ₃	C=C-(2-NO ₂) Ph	H
381	5,6-diF	CF ₃	C=C-(3-NO ₂) Ph	H
382	5,6-diF	CF ₃	C=C-(4-NO ₂) Ph	H
383	5,6-diF	CF ₃	C=C-(2-NH ₂) Ph	H
384	5,6-diF	CF ₃	C=C-(3-NH ₂) Ph	H
385	5,6-diF	CF ₃	C=C-(4-NH ₂) Ph	H
386	5,6-diF	CF ₃	C=C-(2-NMe ₂) Ph	H
387	5,6-diF	CF ₃	C=C-(3-NMe ₂) Ph	H
388	5,6-diF	CF ₃	C=C-(4-NMe ₂) Ph	H
389	5,6-diF	CF ₃	C=C-3-Pyridyl	H
390	5,6-diF	CF ₃	C=C-4-Pyridyl	H
391	5,6-diF	CF ₃	C=C-2-furanyl	H
392	5,6-diF	CF ₃	C=C-3-furanyl	H
393	5,6-diF	CF ₃	C=C-2-thienyl	H
394	5,6-diF	CF ₃	C=C-3-thienyl	H
395	5,6-diF	CF ₃	C=C-2-oxazolyl	H
396	5,6-diF	CF ₃	C=C-2-thiazolyl	H
397	5,6-diF	CF ₃	C=C-4-isoxazolyl	H
398	5,6-diF	CF ₃	C=C-2-imidazolyl	H
399	5,6-diF	CF ₃	CH ₂ CH ₂ -cycPr	H
400	5,6-diF	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ OH	H
401	5,6-diF	CF ₃	CH ₂ CH ₂ -CH(OH)Me	H
402	5,6-diF	CF ₃	CH ₂ CH ₂ -Ph	H

403	5,6-diF	CF ₃	CH ₂ CH ₂ -(2-Cl) Ph	H
404	5,6-diF	CF ₃	CH ₂ CH ₂ -(3-Cl) Ph	H
405	5,6-diF	CF ₃	CH ₂ CH ₂ -(4-Cl) Ph	H
406	5,6-diF	CF ₃	CH ₂ CH ₂ -(2-F) Ph	H
407	5,6-diF	CF ₃	CH ₂ CH ₂ -(3-F) Ph	H
408	5,6-diF	CF ₃	CH ₂ CH ₂ -(4-F) Ph	H
409	5,6-diF	CF ₃	CH ₂ CH ₂ -(2-OH) Ph	H
410	5,6-diF	CF ₃	CH ₂ CH ₂ -(3-OH) Ph	H
411	5,6-diF	CF ₃	CH ₂ CH ₂ -(4-OH) Ph	H
412	5,6-diF	CF ₃	CH ₂ CH ₂ -(2-OMe) Ph	H
413	5,6-diF	CF ₃	CH ₂ CH ₂ -(3-OMe) Ph	H
414	5,6-diF	CF ₃	CH ₂ CH ₂ -(4-OMe) Ph	H
415	5,6-diF	CF ₃	CH ₂ CH ₂ -(2-CN) Ph	H
416	5,6-diF	CF ₃	CH ₂ CH ₂ -(3-CN) Ph	H
417	5,6-diF	CF ₃	CH ₂ CH ₂ -(4-CN) Ph	H
418	5,6-diF	CF ₃	CH ₂ CH ₂ -(2-NO ₂) Ph	H
419	5,6-diF	CF ₃	CH ₂ CH ₂ -(3-NO ₂) Ph	H
420	5,6-diF	CF ₃	CH ₂ CH ₂ -(4-NO ₂) Ph	H
421	5,6-diF	CF ₃	CH ₂ CH ₂ -(2-NH ₂) Ph	H
422	5,6-diF	CF ₃	CH ₂ CH ₂ -(3-NH ₂) Ph	H
423	5,6-diF	CF ₃	CH ₂ CH ₂ -(4-NH ₂) Ph	H
424	5,6-diF	CF ₃	CH ₂ CH ₂ -(2-NMe ₂) Ph	H
425	5,6-diF	CF ₃	CH ₂ CH ₂ -(3-NMe ₂) Ph	H
426	5,6-diF	CF ₃	CH ₂ CH ₂ -(4-NMe ₂) Ph	H
427	5,6-diF	CF ₃	CH ₂ CH ₂ -2-Pyridyl	H
428	5,6-diF	CF ₃	CH ₂ CH ₂ -3-Pyridyl	H
429	5,6-diF	CF ₃	CH ₂ CH ₂ -4-Pyridyl	H
430	5,6-diF	CF ₃	CH ₂ CH ₂ -2-furanyl	H
431	5,6-diF	CF ₃	CH ₂ CH ₂ -3-furanyl	H
432	5,6-diF	CF ₃	CH ₂ CH ₂ -2-thienyl	H
433	5,6-diF	CF ₃	CH ₂ CH ₂ -3-thienyl	H
434	5,6-diF	CF ₃	CH ₂ CH ₂ -2-oxazolyl	H
435	5,6-diF	CF ₃	CH ₂ CH ₂ -2-thiazolyl	H
436	5,6-diF	CF ₃	CH ₂ CH ₂ -4-isoxazolyl	H

437	5,6-diF	CF ₃	CH ₂ CH ₂ -2-imidazolyl	H
438	5,6-diF	CF ₃	C≡C-cycPr	CH ₃
439	5,6-diF	CF ₃	C≡C-2-Pyridyl	CH ₃
440	5,6-diF	CF ₃	C≡C-3-Pyridyl	CH ₃
441	5,6-diF	CF ₃	C≡C-4-Pyridyl	CH ₃
442	5,6-diF	CF ₃	C≡C-2-furanyl	CH ₃
443	5,6-diF	CF ₃	C≡C-3-furanyl	CH ₃
444	5,6-diF	CF ₃	C≡C-2-thienyl	CH ₃
445	5,6-diF	CF ₃	C≡C-3-thienyl	CH ₃
446	5,6-diF	CF ₃	C=C-cycPr	CH ₃
447	5,6-diF	CF ₃	C=C-2-Pyridyl	CH ₃
448	5,6-diF	CF ₃	C=C-3-Pyridyl	CH ₃
449	5,6-diF	CF ₃	C=C-4-Pyridyl	CH ₃
450	5,6-diF	CF ₃	C=C-2-furanyl	CH ₃
451	5,6-diF	CF ₃	C=C-3-furanyl	CH ₃
452	5,6-diF	CF ₃	C=C-2-thienyl	CH ₃
453	5,6-diF	CF ₃	C=C-3-thienyl	CH ₃
454	5,6-diF	CF ₃	CH ₂ CH ₂ -cycPr	CH ₃
455	5,6-diF	CF ₃	CH ₂ CH ₂ -Ph	CH ₃
456	5,6-diF	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₃
457	5,6-diF	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₃
458	5,6-diF	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₃
459	5,6-diF	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₃
460	5,6-diF	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₃
461	5,6-diF	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₃
462	5,6-diF	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₃
463	5,6-diF	CF ₃	C≡C-cycPr	CH ₂ CH ₃
464	5,6-diF	CF ₃	C≡C-Ph	CH ₂ CH ₃
465	5,6-diF	CF ₃	C≡C-2-Pyridyl	CH ₂ CH ₃
466	5,6-diF	CF ₃	C≡C-3-Pyridyl	CH ₂ CH ₃
467	5,6-diF	CF ₃	C≡C-4-Pyridyl	CH ₂ CH ₃
468	5,6-diF	CF ₃	C≡C-2-furanyl	CH ₂ CH ₃
469	5,6-diF	CF ₃	C≡C-3-furanyl	CH ₂ CH ₃
470	5,6-diF	CF ₃	C≡C-2-thienyl	CH ₂ CH ₃

471	5,6-diF	CF ₃	C≡C-3-thienyl	CH ₂ CH ₃
472	5,6-diF	CF ₃	C=C-cycPr	CH ₂ CH ₃
473	5,6-diF	CF ₃	C=C-Ph	CH ₂ CH ₃
474	5,6-diF	CF ₃	C=C-2-Pyridyl	CH ₂ CH ₃
475	5,6-diF	CF ₃	C=C-3-Pyridyl	CH ₂ CH ₃
476	5,6-diF	CF ₃	C=C-4-Pyridyl	CH ₂ CH ₃
477	5,6-diF	CF ₃	C=C-2-furanyl	CH ₂ CH ₃
478	5,6-diF	CF ₃	C=C-3-furanyl	CH ₂ CH ₃
479	5,6-diF	CF ₃	C=C-2-thienyl	CH ₂ CH ₃
480	5,6-diF	CF ₃	C=C-3-thienyl	CH ₂ CH ₃
481	5,6-diF	CF ₃	CH ₂ CH ₂ -cycPr	CH ₂ CH ₃
482	5,6-diF	CF ₃	CH ₂ CH ₂ -Ph	CH ₂ CH ₃
483	5,6-diF	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₂ CH ₃
484	5,6-diF	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₂ CH ₃
485	5,6-diF	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₂ CH ₃
486	5,6-diF	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₂ CH ₃
487	5,6-diF	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₂ CH ₃
488	5,6-diF	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₂ CH ₃
489	5,6-diF	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₂ CH ₃
490	5,6-diCl	CF ₃	C≡C-(2-Cl) Ph	H
491	5,6-diCl	CF ₃	C≡C-(3-Cl) Ph	H
492	5,6-diCl	CF ₃	C≡C-(4-Cl) Ph	H
493	5,6-diCl	CF ₃	C≡C-(2-F) Ph	H
494	5,6-diCl	CF ₃	C≡C-(3-F) Ph	H
495	5,6-diCl	CF ₃	C≡C-(4-F) Ph	H
496	5,6-diCl	CF ₃	C≡C-(2-OH) Ph	H
497	5,6-diCl	CF ₃	C≡C-(3-OH) Ph	H
498	5,6-diCl	CF ₃	C≡C-(4-OH) Ph	H
499	5,6-diCl	CF ₃	C≡C-(2-OMe) Ph	H
500	5,6-diCl	CF ₃	C≡C-(3-OMe) Ph	H
501	5,6-diCl	CF ₃	C≡C-(4-OMe) Ph	H
502	5,6-diCl	CF ₃	C≡C-(2-CN) Ph	H
503	5,6-diCl	CF ₃	C≡C-(3-CN) Ph	H
504	5,6-diCl	CF ₃	C≡C-(4-CN) Ph	H

505	5,6-diCl	CF ₃	C≡C-(2-NO ₂) Ph	H
506	5,6-diCl	CF ₃	C≡C-(3-NO ₂) Ph	H
507	5,6-diCl	CF ₃	C≡C-(4-NO ₂) Ph	H
508	5,6-diCl	CF ₃	C≡C-(2-NH ₂) Ph	H
509	5,6-diCl	CF ₃	C≡C-(3-NH ₂) Ph	H
510	5,6-diCl	CF ₃	C≡C-(4-NH ₂) Ph	H
511	5,6-diCl	CF ₃	C≡C-(2-NMe ₂) Ph	H
512	5,6-diCl	CF ₃	C≡C-(3-NMe ₂) Ph	H
513	5,6-diCl	CF ₃	C≡C-(4-NMe ₂) Ph	H
514	5,6-diCl	CF ₃	C≡C-3-Pyridyl	H
515	5,6-diCl	CF ₃	C≡C-4-Pyridyl	H
516	5,6-diCl	CF ₃	C≡C-2-furanyl	H
517	5,6-diCl	CF ₃	C≡C-3-furanyl	H
518	5,6-diCl	CF ₃	C≡C-2-thienyl	H
519	5,6-diCl	CF ₃	C≡C-3-thienyl	H
520	5,6-diCl	CF ₃	C≡C-2-oxazolyl	H
521	5,6-diCl	CF ₃	C≡C-2-thiazolyl	H
522	5,6-diCl	CF ₃	C≡C-4-isoxazolyl	H
523	5,6-diCl	CF ₃	C≡C-2-imidazolyl	H
524	5,6-diCl	CF ₃	C=C-(2-Cl) Ph	H
525	5,6-diCl	CF ₃	C=C-(3-Cl) Ph	H
526	5,6-diCl	CF ₃	C=C-(4-Cl) Ph	H
527	5,6-diCl	CF ₃	C=C-(2-F) Ph	H
528	5,6-diCl	CF ₃	C=C-(3-F) Ph	H
529	5,6-diCl	CF ₃	C=C-(4-F) Ph	H
530	5,6-diCl	CF ₃	C=C-(2-OH) Ph	H
531	5,6-diCl	CF ₃	C=C-(3-OH) Ph	H
532	5,6-diCl	CF ₃	C=C-(4-OH) Ph	H
533	5,6-diCl	CF ₃	C=C-(2-OMe) Ph	H
534	5,6-diCl	CF ₃	C=C-(3-OMe) Ph	H
535	5,6-diCl	CF ₃	C=C-(4-OMe) Ph	H
536	5,6-diCl	CF ₃	C=C-(2-CN) Ph	H
537	5,6-diCl	CF ₃	C=C-(3-CN) Ph	H
538	5,6-diCl	CF ₃	C=C-(4-CN) Ph	H

539	5,6-diCl	CF ₃	C=C-(2-NO ₂) Ph	H
540	5,6-diCl	CF ₃	C=C-(3-NO ₂) Ph	H
541	5,6-diCl	CF ₃	C=C-(4-NO ₂) Ph	H
542	5,6-diCl	CF ₃	C=C-(2-NH ₂) Ph	H
543	5,6-diCl	CF ₃	C=C-(3-NH ₂) Ph	H
544	5,6-diCl	CF ₃	C=C-(4-NH ₂) Ph	H
545	5,6-diCl	CF ₃	C=C-(2-NMe ₂) Ph	H
546	5,6-diCl	CF ₃	C=C-(3-NMe ₂) Ph	H
547	5,6-diCl	CF ₃	C=C-(4-NMe ₂) Ph	H
548	5,6-diCl	CF ₃	C=C-3-Pyridyl	H
549	5,6-diCl	CF ₃	C=C-4-Pyridyl	H
550	5,6-diCl	CF ₃	C=C-2-furanyl	H
551	5,6-diCl	CF ₃	C=C-3-furanyl	H
552	5,6-diCl	CF ₃	C=C-2-thienyl	H
553	5,6-diCl	CF ₃	C=C-3-thienyl	H
554	5,6-diCl	CF ₃	C=C-2-oxazolyl	H
555	5,6-diCl	CF ₃	C=C-2-thiazolyl	H
556	5,6-diCl	CF ₃	C=C-4-isoxazolyl	H
557	5,6-diCl	CF ₃	C=C-2-imidazolyl	H
558	5,6-diCl	CF ₃	CH ₂ CH ₂ -cycPr	H
559	5,6-diCl	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ OH	H
560	5,6-diCl	CF ₃	CH ₂ CH ₂ -CH(OH)Me	H
561	5,6-diCl	CF ₃	CH ₂ CH ₂ -Ph	H
562	5,6-diCl	CF ₃	CH ₂ CH ₂ -(2-Cl) Ph	H
563	5,6-diCl	CF ₃	CH ₂ CH ₂ -(3-Cl) Ph	H
564	5,6-diCl	CF ₃	CH ₂ CH ₂ -(4-Cl) Ph	H
565	5,6-diCl	CF ₃	CH ₂ CH ₂ -(2-F) Ph	H
566	5,6-diCl	CF ₃	CH ₂ CH ₂ -(3-F) Ph	H
567	5,6-diCl	CF ₃	CH ₂ CH ₂ -(4-F) Ph	H
568	5,6-diCl	CF ₃	CH ₂ CH ₂ -(2-OH) Ph	H
569	5,6-diCl	CF ₃	CH ₂ CH ₂ -(3-OH) Ph	H
570	5,6-diCl	CF ₃	CH ₂ CH ₂ -(4-OH) Ph	H
571	5,6-diCl	CF ₃	CH ₂ CH ₂ -(2-OMe) Ph	H
572	5,6-diCl	CF ₃	CH ₂ CH ₂ -(3-OMe) Ph	H

573	5,6-diCl	CF ₃	CH ₂ CH ₂ -(4-OMe) Ph	H
574	5,6-diCl	CF ₃	CH ₂ CH ₂ -(2-CN) Ph	H
575	5,6-diCl	CF ₃	CH ₂ CH ₂ -(3-CN) Ph	H
576	5,6-diCl	CF ₃	CH ₂ CH ₂ -(4-CN) Ph	H
577	5,6-diCl	CF ₃	CH ₂ CH ₂ -(2-NO ₂) Ph	H
578	5,6-diCl	CF ₃	CH ₂ CH ₂ -(3-NO ₂) Ph	H
579	5,6-diCl	CF ₃	CH ₂ CH ₂ -(4-NO ₂) Ph	H
580	5,6-diCl	CF ₃	CH ₂ CH ₂ -(2-NH ₂) Ph	H
581	5,6-diCl	CF ₃	CH ₂ CH ₂ -(3-NH ₂) Ph	H
582	5,6-diCl	CF ₃	CH ₂ CH ₂ -(4-NH ₂) Ph	H
583	5,6-diCl	CF ₃	CH ₂ CH ₂ -(2-NMe ₂) Ph	H
584	5,6-diCl	CF ₃	CH ₂ CH ₂ -(3-NMe ₂) Ph	H
585	5,6-diCl	CF ₃	CH ₂ CH ₂ -(4-NMe ₂) Ph	H
586	5,6-diCl	CF ₃	CH ₂ CH ₂ -2-Pyridyl	H
587	5,6-diCl	CF ₃	CH ₂ CH ₂ -3-Pyridyl	H
588	5,6-diCl	CF ₃	CH ₂ CH ₂ -4-Pyridyl	H
589	5,6-diCl	CF ₃	CH ₂ CH ₂ -2-furanyl	H
590	5,6-diCl	CF ₃	CH ₂ CH ₂ -3-furanyl	H
591	5,6-diCl	CF ₃	CH ₂ CH ₂ -2-thienyl	H
592	5,6-diCl	CF ₃	CH ₂ CH ₂ -3-thienyl	H
593	5,6-diCl	CF ₃	CH ₂ CH ₂ -2-oxazolyl	H
594	5,6-diCl	CF ₃	CH ₂ CH ₂ -2-thiazolyl	H
595	5,6-diCl	CF ₃	CH ₂ CH ₂ -4-isoxazolyl	H
596	5,6-diCl	CF ₃	CH ₂ CH ₂ -2-imidazolyl	H
597	5,6-diCl	CF ₃	C≡C-cycPr	CH ₃
598	5,6-diCl	CF ₃	C≡C-2-Pyridyl	CH ₃
599	5,6-diCl	CF ₃	C≡C-3-Pyridyl	CH ₃
600	5,6-diCl	CF ₃	C≡C-4-Pyridyl	CH ₃
601	5,6-diCl	CF ₃	C≡C-2-furanyl	CH ₃
602	5,6-diCl	CF ₃	C≡C-3-furanyl	CH ₃
603	5,6-diCl	CF ₃	C≡C-2-thienyl	CH ₃
604	5,6-diCl	CF ₃	C≡C-3-thienyl	CH ₃
605	5,6-diCl	CF ₃	C=C-cycPr	CH ₃
606	5,6-diCl	CF ₃	C=C-2-Pyridyl	CH ₃

607	5,6-diCl	CF ₃	C=C-3-Pyridyl	CH ₃
608	5,6-diCl	CF ₃	C=C-4-Pyridyl	CH ₃
609	5,6-diCl	CF ₃	C=C-2-furanyl	CH ₃
610	5,6-diCl	CF ₃	C=C-3-furanyl	CH ₃
611	5,6-diCl	CF ₃	C=C-2-thienyl	CH ₃
612	5,6-diCl	CF ₃	C=C-3-thienyl	CH ₃
613	5,6-diCl	CF ₃	CH ₂ CH ₂ -cycPr	CH ₃
614	5,6-diCl	CF ₃	CH ₂ CH ₂ -Ph	CH ₃
615	5,6-diCl	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₃
616	5,6-diCl	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₃
617	5,6-diCl	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₃
618	5,6-diCl	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₃
619	5,6-diCl	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₃
620	5,6-diCl	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₃
621	5,6-diCl	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₃
622	5,6-diCl	CF ₃	C≡C-cycPr	CH ₂ CH ₃
623	5,6-diCl	CF ₃	C≡C-Ph	CH ₂ CH ₃
624	5,6-diCl	CF ₃	C≡C-2-Pyridyl	CH ₂ CH ₃
625	5,6-diCl	CF ₃	C≡C-3-Pyridyl	CH ₂ CH ₃
626	5,6-diCl	CF ₃	C≡C-4-Pyridyl	CH ₂ CH ₃
627	5,6-diCl	CF ₃	C≡C-2-furanyl	CH ₂ CH ₃
628	5,6-diCl	CF ₃	C≡C-3-furanyl	CH ₂ CH ₃
629	5,6-diCl	CF ₃	C≡C-2-thienyl	CH ₂ CH ₃
630	5,6-diCl	CF ₃	C≡C-3-thienyl	CH ₂ CH ₃
631	5,6-diCl	CF ₃	C=C-cycPr	CH ₂ CH ₃
632	5,6-diCl	CF ₃	C=C-Ph	CH ₂ CH ₃
633	5,6-diCl	CF ₃	C=C-2-Pyridyl	CH ₂ CH ₃
634	5,6-diCl	CF ₃	C=C-3-Pyridyl	CH ₂ CH ₃
635	5,6-diCl	CF ₃	C=C-4-Pyridyl	CH ₂ CH ₃
636	5,6-diCl	CF ₃	C=C-2-furanyl	CH ₂ CH ₃
637	5,6-diCl	CF ₃	C=C-3-furanyl	CH ₂ CH ₃
638	5,6-diCl	CF ₃	C=C-2-thienyl	CH ₂ CH ₃
639	5,6-diCl	CF ₃	C=C-3-thienyl	CH ₂ CH ₃
640	5,6-diCl	CF ₃	CH ₂ CH ₂ -cycPr	CH ₂ CH ₃

641	5,6-diCl	CF ₃	CH ₂ CH ₂ -Ph	CH ₂ CH ₃
642	5,6-diCl	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₂ CH ₃
643	5,6-diCl	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₂ CH ₃
644	5,6-diCl	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₂ CH ₃
645	5,6-diCl	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₂ CH ₃
646	5,6-diCl	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₂ CH ₃
647	5,6-diCl	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₂ CH ₃
648	5,6-diCl	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₂ CH ₃
649	6-F	CF ₃	C≡CCH ₂ CH ₂ OH	H
650	6-F	CF ₃	C≡C-CH(OH)Me	H
651	6-F	CF ₃	C≡C-(2-Cl)Ph	H
652	6-F	CF ₃	C≡C-(3-Cl)Ph	H
653	6-F	CF ₃	C≡C-(4-Cl)Ph	H
654	6-F	CF ₃	C≡C-(2-F)Ph	H
655	6-F	CF ₃	C≡C-(3-F)Ph	H
656	6-F	CF ₃	C≡C-(4-F)Ph	H
657	6-F	CF ₃	C≡C-(2-OH)Ph	H
658	6-F	CF ₃	C≡C-(3-OH)Ph	H
659	6-F	CF ₃	C≡C-(4-OH)Ph	H
660	6-F	CF ₃	C≡C-(2-OMe)Ph	H
661	6-F	CF ₃	C≡C-(3-OMe)Ph	H
662	6-F	CF ₃	C≡C-(4-OMe)Ph	H
663	6-F	CF ₃	C≡C-(2-CN)Ph	H
664	6-F	CF ₃	C≡C-(3-CN)Ph	H
665	6-F	CF ₃	C≡C-(4-CN)Ph	H
666	6-F	CF ₃	C≡C-(2-NO ₂)Ph	H
667	6-F	CF ₃	C≡C-(3-NO ₂)Ph	H
668	6-F	CF ₃	C≡C-(4-NO ₂)Ph	H
669	6-F	CF ₃	C≡C-(2-NH ₂)Ph	H
670	6-F	CF ₃	C≡C-(3-NH ₂)Ph	H
671	6-F	CF ₃	C≡C-(4-NH ₂)Ph	H
672	6-F	CF ₃	C≡C-(2-NMe ₂)Ph	H
673	6-F	CF ₃	C≡C-(3-NMe ₂)Ph	H
674	6-F	CF ₃	C≡C-(4-NMe ₂)Ph	H

675	6-F	CF ₃	C≡C-3-Pyridyl	H
676	6-F	CF ₃	C≡C-4-Pyridyl	H
677	6-F	CF ₃	C≡C-2-furanyl	H
678	6-F	CF ₃	C≡C-3-furanyl	H
679	6-F	CF ₃	C≡C-2-thienyl	H
680	6-F	CF ₃	C≡C-3-thienyl	H
681	6-F	CF ₃	C≡C-2-oxazolyl	H
682	6-F	CF ₃	C≡C-2-thiazolyl	H
683	6-F	CF ₃	C≡C-4-isoxazolyl	H
684	6-F	CF ₃	C≡C-2-imidazolyl	H
685	6-F	CF ₃	C=CCH ₂ CH ₂ OH	H
686	6-F	CF ₃	C=C-CH(OH)Me	H
687	6-F	CF ₃	C=C-(2-Cl)Ph	H
688	6-F	CF ₃	C=C-(3-Cl)Ph	H
689	6-F	CF ₃	C=C-(4-Cl)Ph	H
690	6-F	CF ₃	C=C-(2-F)Ph	H
691	6-F	CF ₃	C=C-(3-F)Ph	H
692	6-F	CF ₃	C=C-(4-F)Ph	H
693	6-F	CF ₃	C=C-(2-OH)Ph	H
694	6-F	CF ₃	C=C-(3-OH)Ph	H
695	6-F	CF ₃	C=C-(4-OH)Ph	H
696	6-F	CF ₃	C=C-(2-OMe)Ph	H
697	6-F	CF ₃	C=C-(3-OMe)Ph	H
698	6-F	CF ₃	C=C-(4-OMe)Ph	H
699	6-F	CF ₃	C=C-(2-CN)Ph	H
700	6-F	CF ₃	C=C-(3-CN)Ph	H
701	6-F	CF ₃	C=C-(4-CN)Ph	H
702	6-F	CF ₃	C=C-(2-NO ₂)Ph	H
703	6-F	CF ₃	C=C-(3-NO ₂)Ph	H
704	6-F	CF ₃	C=C-(4-NO ₂)Ph	H
705	6-F	CF ₃	C=C-(2-NH ₂)Ph	H
706	6-F	CF ₃	C=C-(3-NH ₂)Ph	H
707	6-F	CF ₃	C=C-(4-NH ₂)Ph	H
708	6-F	CF ₃	C=C-(2-NMe ₂)Ph	H

709	6-F	CF ₃	C=C- (3-NMe ₂) Ph	H
710	6-F	CF ₃	C=C- (4-NMe ₂) Ph	H
711	6-F	CF ₃	C=C-3-Pyridyl	H
712	6-F	CF ₃	C=C-4-Pyridyl	H
713	6-F	CF ₃	C=C-2-furanyl	H
714	6-F	CF ₃	C=C-3-furanyl	H
715	6-F	CF ₃	C=C-2-thienyl	H
716	6-F	CF ₃	C=C-3-thienyl	H
717	6-F	CF ₃	C=C-2-oxazolyl	H
718	6-F	CF ₃	C=C-2-thiazolyl	H
719	6-F	CF ₃	C=C-4-isoxazolyl	H
720	6-F	CF ₃	C=C-2-imidazolyl	H
721	6-F	CF ₃	CH ₂ CH ₂ -cycPr	H
722	6-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ OH	H
723	6-F	CF ₃	CH ₂ CH ₂ -CH(OH)Me	H
724	6-F	CF ₃	CH ₂ CH ₂ - (2-Cl) Ph	H
725	6-F	CF ₃	CH ₂ CH ₂ - (3-Cl) Ph	H
726	6-F	CF ₃	CH ₂ CH ₂ - (4-Cl) Ph	H
727	6-F	CF ₃	CH ₂ CH ₂ - (2-F) Ph	H
728	6-F	CF ₃	CH ₂ CH ₂ - (3-F) Ph	H
729	6-F	CF ₃	CH ₂ CH ₂ - (4-F) Ph	H
730	6-F	CF ₃	CH ₂ CH ₂ - (2-OH) Ph	H
731	6-F	CF ₃	CH ₂ CH ₂ - (3-OH) Ph	H
732	6-F	CF ₃	CH ₂ CH ₂ - (4-OH) Ph	H
733	6-F	CF ₃	CH ₂ CH ₂ - (2-OMe) Ph	H
734	6-F	CF ₃	CH ₂ CH ₂ - (3-OMe) Ph	H
735	6-F	CF ₃	CH ₂ CH ₂ - (4-OMe) Ph	H
736	6-F	CF ₃	CH ₂ CH ₂ - (2-CN) Ph	H
737	6-F	CF ₃	CH ₂ CH ₂ - (3-CN) Ph	H
738	6-F	CF ₃	CH ₂ CH ₂ - (4-CN) Ph	H
739	6-F	CF ₃	CH ₂ CH ₂ - (2-NO ₂) Ph	H
740	6-F	CF ₃	CH ₂ CH ₂ - (3-NO ₂) Ph	H
741	6-F	CF ₃	CH ₂ CH ₂ - (4-NO ₂) Ph	H
742	6-F	CF ₃	CH ₂ CH ₂ - (2-NH ₂) Ph	H

743	6-F	CF ₃	CH ₂ CH ₂ -(3-NH ₂) Ph	H
744	6-F	CF ₃	CH ₂ CH ₂ -(4-NH ₂) Ph	H
745	6-F	CF ₃	CH ₂ CH ₂ -(2-NMe ₂) Ph	H
746	6-F	CF ₃	CH ₂ CH ₂ -(3-NMe ₂) Ph	H
747	6-F	CF ₃	CH ₂ CH ₂ -(4-NMe ₂) Ph	H
748	6-F	CF ₃	CH ₂ CH ₂ -3-Pyridyl	H
749	6-F	CF ₃	CH ₂ CH ₂ -4-Pyridyl	H
750	6-F	CF ₃	CH ₂ CH ₂ -2-furanyl	H
751	6-F	CF ₃	CH ₂ CH ₂ -3-furanyl	H
752	6-F	CF ₃	CH ₂ CH ₂ -2-thienyl	H
753	6-F	CF ₃	CH ₂ CH ₂ -3-thienyl	H
754	6-F	CF ₃	CH ₂ CH ₂ -2-oxazolyl	H
755	6-F	CF ₃	CH ₂ CH ₂ -2-thiazolyl	H
756	6-F	CF ₃	CH ₂ CH ₂ -4-isoxazolyl	H
757	6-F	CF ₃	CH ₂ CH ₂ -2-imidazolyl	H
758	6-F	CF ₃	C≡C-cycPr	CH ₃
759	6-F	CF ₃	C≡C-iPr	CH ₃
760	6-F	CF ₃	C≡C-Pr	CH ₃
761	6-F	CF ₃	C≡C-Bu	CH ₃
762	6-F	CF ₃	C≡C-iBu	CH ₃
763	6-F	CF ₃	C≡C-tBu	CH ₃
764	6-F	CF ₃	C≡C-Et	CH ₃
765	6-F	CF ₃	C≡C-Me	CH ₃
766	6-F	CF ₃	C≡C-Ph	CH ₃
767	6-F	CF ₃	C≡C-2-Pyridyl	CH ₃
768	6-F	CF ₃	C≡C-3-Pyridyl	CH ₃
769	6-F	CF ₃	C≡C-4-Pyridyl	CH ₃
770	6-F	CF ₃	C≡C-2-furanyl	CH ₃
771	6-F	CF ₃	C≡C-3-furanyl	CH ₃
772	6-F	CF ₃	C≡C-2-thienyl	CH ₃
773	6-F	CF ₃	C≡C-3-thienyl	CH ₃
774	6-F	CF ₃	C=C-cycPr	CH ₃
775	6-F	CF ₃	C=C-iPr	CH ₃
776	6-F	CF ₃	C=C-Pr	CH ₃

777	6-F	CF ₃	C=C-Bu	CH ₃
778	6-F	CF ₃	C=C-iBu	CH ₃
779	6-F	CF ₃	C=C-tBu	CH ₃
780	6-F	CF ₃	C=C-Et	CH ₃
781	6-F	CF ₃	C=C-Me	CH ₃
782	6-F	CF ₃	C=C-Ph	CH ₃
783	6-F	CF ₃	C=C-2-Pyridyl	CH ₃
784	6-F	CF ₃	C=C-3-Pyridyl	CH ₃
785	6-F	CF ₃	C=C-4-Pyridyl	CH ₃
786	6-F	CF ₃	C=C-2-furanyl	CH ₃
787	6-F	CF ₃	C=C-3-furanyl	CH ₃
788	6-F	CF ₃	C=C-2-thienyl	CH ₃
789	6-F	CF ₃	C=C-3-thienyl	CH ₃
790	6-F	CF ₃	CH ₂ CH ₂ -cycPr	CH ₃
791	6-F	CF ₃	CH ₂ CH ₂ -Ph	CH ₃
792	6-F	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₃
793	6-F	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₃
794	6-F	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₃
795	6-F	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₃
796	6-F	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₃
797	6-F	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₃
798	6-F	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₃
799	6-F	CF ₃	C≡C-cycPr	CH ₂ CH ₃
800	6-F	CF ₃	C≡C-Ph	CH ₂ CH ₃
801	6-F	CF ₃	C≡C-2-Pyridyl	CH ₂ CH ₃
802	6-F	CF ₃	C≡C-3-Pyridyl	CH ₂ CH ₃
803	6-F	CF ₃	C≡C-4-Pyridyl	CH ₂ CH ₃
804	6-F	CF ₃	C≡C-2-furanyl	CH ₂ CH ₃
805	6-F	CF ₃	C≡C-3-furanyl	CH ₂ CH ₃
806	6-F	CF ₃	C≡C-2-thienyl	CH ₂ CH ₃
807	6-F	CF ₃	C≡C-3-thienyl	CH ₂ CH ₃
808	6-F	CF ₃	C=C-cycPr	CH ₂ CH ₃
809	6-F	CF ₃	C=C-Ph	CH ₂ CH ₃
810	6-F	CF ₃	C=C-2-Pyridyl	CH ₂ CH ₃

811	6-F	CF ₃	C=C-3-Pyridyl	CH ₂ CH ₃
812	6-F	CF ₃	C=C-4-Pyridyl	CH ₂ CH ₃
813	6-F	CF ₃	C=C-2-furanyl	CH ₂ CH ₃
814	6-F	CF ₃	C=C-3-furanyl	CH ₂ CH ₃
815	6-F	CF ₃	C=C-2-thienyl	CH ₂ CH ₃
816	6-F	CF ₃	C=C-3-thienyl	CH ₂ CH ₃
817	6-F	CF ₃	CH ₂ CH ₂ -cycPr	CH ₂ CH ₃
818	6-F	CF ₃	CH ₂ CH ₂ -Ph	CH ₂ CH ₃
819	6-F	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₂ CH ₃
820	6-F	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₂ CH ₃
821	6-F	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₂ CH ₃
822	6-F	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₂ CH ₃
823	6-F	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₂ CH ₃
824	6-F	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₂ CH ₃
825	6-F	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₂ CH ₃
826	5-Cl	CF ₃	C≡C-cycPr	H
827	5-Cl	CF ₃	C≡CCH ₂ CH ₂ OH	H
828	5-Cl	CF ₃	C≡C-CH(OH)Me	H
829	5-Cl	CF ₃	C≡C-Ph	H
830	5-Cl	CF ₃	C≡C-(2-Cl)Ph	H
831	5-Cl	CF ₃	C≡C-(3-Cl)Ph	H
832	5-Cl	CF ₃	C≡C-(4-Cl)Ph	H
833	5-Cl	CF ₃	C≡C-(2-F)Ph	H
834	5-Cl	CF ₃	C≡C-(3-F)Ph	H
835	5-Cl	CF ₃	C≡C-(4-F)Ph	H
836	5-Cl	CF ₃	C≡C-(2-OH)Ph	H
837	5-Cl	CF ₃	C≡C-(3-OH)Ph	H
838	5-Cl	CF ₃	C≡C-(4-OH)Ph	H
839	5-Cl	CF ₃	C≡C-(2-OMe)Ph	H
840	5-Cl	CF ₃	C≡C-(3-OMe)Ph	H
841	5-Cl	CF ₃	C≡C-(4-OMe)Ph	H
842	5-Cl	CF ₃	C≡C-(2-CN)Ph	H
843	5-Cl	CF ₃	C≡C-(3-CN)Ph	H
844	5-Cl	CF ₃	C≡C-(4-CN)Ph	H

845	5-Cl	CF ₃	C≡C-(2-NO ₂) Ph	H
846	5-Cl	CF ₃	C≡C-(3-NO ₂) Ph	H
847	5-Cl	CF ₃	C≡C-(4-NO ₂) Ph	H
848	5-Cl	CF ₃	C≡C-(2-NH ₂) Ph	H
849	5-Cl	CF ₃	C≡C-(3-NH ₂) Ph	H
850	5-Cl	CF ₃	C≡C-(4-NH ₂) Ph	H
851	5-Cl	CF ₃	C≡C-(2-NMe ₂) Ph	H
852	5-Cl	CF ₃	C≡C-(3-NMe ₂) Ph	H
853	5-Cl	CF ₃	C≡C-(4-NMe ₂) Ph	H
854	5-Cl	CF ₃	C≡C-2-Pyridyl	H
855	5-Cl	CF ₃	C≡C-2-Pyridyl	H
856	5-Cl	CF ₃	C≡C-3-Pyridyl	H
857	5-Cl	CF ₃	C≡C-4-Pyridyl	H
858	5-Cl	CF ₃	C≡C-2-furanyl	H
859	5-Cl	CF ₃	C≡C-3-furanyl	H
860	5-Cl	CF ₃	C≡C-2-thienyl	H
861	5-Cl	CF ₃	C≡C-3-thienyl	H
862	5-Cl	CF ₃	C≡C-2-oxazolyl	H
863	5-Cl	CF ₃	C≡C-2-thiazolyl	H
864	5-Cl	CF ₃	C≡C-4-isoxazolyl	H
865	5-Cl	CF ₃	C≡C-2-imidazolyl	H
866	5-Cl	CF ₃	C=C-cycPr	H
867	5-Cl	CF ₃	C=CCH ₂ CH ₂ OH	H
868	5-Cl	CF ₃	C=C-CH(OH)Me	H
869	5-Cl	CF ₃	C=C-Ph	H
870	5-Cl	CF ₃	C=C-(2-Cl) Ph	H
871	5-Cl	CF ₃	C=C-(3-Cl) Ph	H
872	5-Cl	CF ₃	C=C-(4-Cl) Ph	H
873	5-Cl	CF ₃	C=C-(2-F) Ph	H
874	5-Cl	CF ₃	C=C-(3-F) Ph	H
875	5-Cl	CF ₃	C=C-(4-F) Ph	H
876	5-Cl	CF ₃	C=C-(2-OH) Ph	H
877	5-Cl	CF ₃	C=C-(3-OH) Ph	H
878	5-Cl	CF ₃	C=C-(4-OH) Ph	H

879	5-Cl	CF ₃	C=C-(2-OMe) Ph	H
880	5-Cl	CF ₃	C=C-(3-OMe) Ph	H
881	5-Cl	CF ₃	C=C-(4-OMe) Ph	H
882	5-Cl	CF ₃	C=C-(2-CN) Ph	H
883	5-Cl	CF ₃	C=C-(3-CN) Ph	H
884	5-Cl	CF ₃	C=C-(4-CN) Ph	H
885	5-Cl	CF ₃	C=C-(2-NO ₂) Ph	H
886	5-Cl	CF ₃	C=C-(3-NO ₂) Ph	H
887	5-Cl	CF ₃	C=C-(4-NO ₂) Ph	H
888	5-Cl	CF ₃	C=C-(2-NH ₂) Ph	H
889	5-Cl	CF ₃	C=C-(3-NH ₂) Ph	H
890	5-Cl	CF ₃	C=C-(4-NH ₂) Ph	H
891	5-Cl	CF ₃	C=C-(2-NMe ₂) Ph	H
892	5-Cl	CF ₃	C=C-(3-NMe ₂) Ph	H
893	5-Cl	CF ₃	C=C-(4-NMe ₂) Ph	H
894	5-Cl	CF ₃	C=C-2-Pyridyl	H
895	5-Cl	CF ₃	C=C-2-Pyridyl	H
896	5-Cl	CF ₃	C=C-3-Pyridyl	H
897	5-Cl	CF ₃	C=C-4-Pyridyl	H
898	5-Cl	CF ₃	C=C-2-furanyl	H
899	5-Cl	CF ₃	C=C-3-furanyl	H
900	5-Cl	CF ₃	C=C-2-thienyl	H
901	5-Cl	CF ₃	C=C-3-thienyl	H
902	5-Cl	CF ₃	C=C-2-oxazolyl	H
903	5-Cl	CF ₃	C=C-2-thiazolyl	H
904	5-Cl	CF ₃	C=C-4-isoxazolyl	H
905	5-Cl	CF ₃	C=C-2-imidazolyl	H
906	5-Cl	CF ₃	CH ₂ CH ₂ -cycPr	H
907	5-Cl	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ OH	H
908	5-Cl	CF ₃	CH ₂ CH ₂ -CH(OH) Me	H
909	5-Cl	CF ₃	CH ₂ CH ₂ Ph	H
910	5-Cl	CF ₃	CH ₂ CH ₂ -(2-Cl) Ph	H
911	5-Cl	CF ₃	CH ₂ CH ₂ -(3-Cl) Ph	H
912	5-Cl	CF ₃	CH ₂ CH ₂ -(4-Cl) Ph	H

913	5-Cl	CF ₃	CH ₂ CH ₂ -(2-F) Ph	H
914	5-Cl	CF ₃	CH ₂ CH ₂ -(3-F) Ph	H
915	5-Cl	CF ₃	CH ₂ CH ₂ -(4-F) Ph	H
916	5-Cl	CF ₃	CH ₂ CH ₂ -(2-OH) Ph	H
917	5-Cl	CF ₃	CH ₂ CH ₂ -(3-OH) Ph	H
918	5-Cl	CF ₃	CH ₂ CH ₂ -(4-OH) Ph	H
919	5-Cl	CF ₃	CH ₂ CH ₂ -(2-OMe) Ph	H
920	5-Cl	CF ₃	CH ₂ CH ₂ -(3-OMe) Ph	H
921	5-Cl	CF ₃	CH ₂ CH ₂ -(4-OMe) Ph	H
922	5-Cl	CF ₃	CH ₂ CH ₂ -(2-CN) Ph	H
923	5-Cl	CF ₃	CH ₂ CH ₂ -(3-CN) Ph	H
924	5-Cl	CF ₃	CH ₂ CH ₂ -(4-CN) Ph	H
925	5-Cl	CF ₃	CH ₂ CH ₂ -(2-NO ₂) Ph	H
926	5-Cl	CF ₃	CH ₂ CH ₂ -(3-NO ₂) Ph	H
927	5-Cl	CF ₃	CH ₂ CH ₂ -(4-NO ₂) Ph	H
928	5-Cl	CF ₃	CH ₂ CH ₂ -(2-NH ₂) Ph	H
929	5-Cl	CF ₃	CH ₂ CH ₂ -(3-NH ₂) Ph	H
930	5-Cl	CF ₃	CH ₂ CH ₂ -(4-NH ₂) Ph	H
931	5-Cl	CF ₃	CH ₂ CH ₂ -(2-NMe ₂) Ph	H
932	5-Cl	CF ₃	CH ₂ CH ₂ -(3-NMe ₂) Ph	H
933	5-Cl	CF ₃	CH ₂ CH ₂ -(4-NMe ₂) Ph	H
934	5-Cl	CF ₃	CH ₂ CH ₂ -2-Pyridyl	H
935	5-Cl	CF ₃	CH ₂ CH ₂ -3-Pyridyl	H
936	5-Cl	CF ₃	CH ₂ CH ₂ -4-Pyridyl	H
937	5-Cl	CF ₃	CH ₂ CH ₂ -2-furanyl	H
938	5-Cl	CF ₃	CH ₂ CH ₂ -3-furanyl	H
939	5-Cl	CF ₃	CH ₂ CH ₂ -2-thienyl	H
940	5-Cl	CF ₃	CH ₂ CH ₂ -3-thienyl	H
941	5-Cl	CF ₃	CH ₂ CH ₂ -2-oxazolyl	H
942	5-Cl	CF ₃	CH ₂ CH ₂ -2-thiazolyl	H
943	5-Cl	CF ₃	CH ₂ CH ₂ -4-isoxazolyl	H
944	5-Cl	CF ₃	CH ₂ CH ₂ -2-imidazolyl	H
945	5-Cl	CF ₃	C≡C-cycPr	CH ₃
946	5-Cl	CF ₃	C≡C-Ph	CH ₃

947	5-Cl	CF ₃	C≡C-2-Pyridyl	CH ₃
948	5-Cl	CF ₃	C≡C-3-Pyridyl	CH ₃
949	5-Cl	CF ₃	C≡C-4-Pyridyl	CH ₃
950	5-Cl	CF ₃	C≡C-2-furanyl	CH ₃
951	5-Cl	CF ₃	C≡C-3-furanyl	CH ₃
952	5-Cl	CF ₃	C≡C-2-thienyl	CH ₃
953	5-Cl	CF ₃	C≡C-3-thienyl	CH ₃
954	5-Cl	CF ₃	C=C-cycPr	CH ₃
955	5-Cl	CF ₃	C=C-Ph	CH ₃
956	5-Cl	CF ₃	C=C-2-Pyridyl	CH ₃
957	5-Cl	CF ₃	C=C-3-Pyridyl	CH ₃
958	5-Cl	CF ₃	C=C-4-Pyridyl	CH ₃
959	5-Cl	CF ₃	C=C-2-furanyl	CH ₃
960	5-Cl	CF ₃	C=C-3-furanyl	CH ₃
961	5-Cl	CF ₃	C=C-2-thienyl	CH ₃
962	5-Cl	CF ₃	C=C-3-thienyl	CH ₃
963	5-Cl	CF ₃	CH ₂ CH ₂ -cycPr	CH ₃
964	5-Cl	CF ₃	CH ₂ CH ₂ -Ph	CH ₃
965	5-Cl	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₃
966	5-Cl	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₃
967	5-Cl	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₃
968	5-Cl	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₃
969	5-Cl	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₃
970	5-Cl	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₃
971	5-Cl	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₃
972	5-Cl	CF ₃	C≡C-cycPr	CH ₂ CH ₃
973	5-Cl	CF ₃	C≡C-Ph	CH ₂ CH ₃
974	5-Cl	CF ₃	C≡C-2-Pyridyl	CH ₂ CH ₃
975	5-Cl	CF ₃	C≡C-3-Pyridyl	CH ₂ CH ₃
976	5-Cl	CF ₃	C≡C-4-Pyridyl	CH ₂ CH ₃
977	5-Cl	CF ₃	C≡C-2-furanyl	CH ₂ CH ₃
978	5-Cl	CF ₃	C≡C-3-furanyl	CH ₂ CH ₃
979	5-Cl	CF ₃	C≡C-2-thienyl	CH ₂ CH ₃
980	5-Cl	CF ₃	C≡C-3-thienyl	CH ₂ CH ₃

981	5-Cl	CF ₃	C=C-cycPr	CH ₂ CH ₃
982	5-Cl	CF ₃	C=C-Ph	CH ₂ CH ₃
983	5-Cl	CF ₃	C=C-2-Pyridyl	CH ₂ CH ₃
984	5-Cl	CF ₃	C=C-3-Pyridyl	CH ₂ CH ₃
985	5-Cl	CF ₃	C=C-4-Pyridyl	CH ₂ CH ₃
986	5-Cl	CF ₃	C=C-2-furanyl	CH ₂ CH ₃
987	5-Cl	CF ₃	C=C-3-furanyl	CH ₂ CH ₃
988	5-Cl	CF ₃	C=C-2-thienyl	CH ₂ CH ₃
989	5-Cl	CF ₃	C=C-3-thienyl	CH ₂ CH ₃
990	5-Cl	CF ₃	CH ₂ CH ₂ -cycPr	CH ₂ CH ₃
991	5-Cl	CF ₃	CH ₂ CH ₂ -Ph	CH ₂ CH ₃
992	5-Cl	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₂ CH ₃
993	5-Cl	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₂ CH ₃
994	5-Cl	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₂ CH ₃
995	5-Cl	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₂ CH ₃
996	5-Cl	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₂ CH ₃
997	5-Cl	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₂ CH ₃
998	5-Cl	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₂ CH ₃
999	5-F	CF ₃	C≡C-cycPr	H
1000	5-F	CF ₃	C≡CCH ₂ CH ₂ OH	H
1001	5-F	CF ₃	C≡C-CH(OH)Me	H
1002	5-F	CF ₃	C≡C-Ph	H
1003	5-F	CF ₃	C≡C-(2-Cl)Ph	H
1004	5-F	CF ₃	C≡C-(3-Cl)Ph	H
1005	5-F	CF ₃	C≡C-(4-Cl)Ph	H
1006	5-F	CF ₃	C≡C-(2-F)Ph	H
1007	5-F	CF ₃	C≡C-(3-F)Ph	H
1008	5-F	CF ₃	C≡C-(4-F)Ph	H
1009	5-F	CF ₃	C≡C-(2-OH)Ph	H
1010	5-F	CF ₃	C≡C-(3-OH)Ph	H
1011	5-F	CF ₃	C≡C-(4-OH)Ph	H
1012	5-F	CF ₃	C≡C-(2-OMe)Ph	H
1013	5-F	CF ₃	C≡C-(3-OMe)Ph	H
1014	5-F	CF ₃	C≡C-(4-OMe)Ph	H

1015	5-F	CF ₃	C≡C-(2-CN) Ph	H
1016	5-F	CF ₃	C≡C-(3-CN) Ph	H
1017	5-F	CF ₃	C≡C-(4-CN) Ph	H
1018	5-F	CF ₃	C≡C-(2-NO ₂) Ph	H
1019	5-F	CF ₃	C≡C-(3-NO ₂) Ph	H
1020	5-F	CF ₃	C≡C-(4-NO ₂) Ph	H
1021	5-F	CF ₃	C≡C-(2-NH ₂) Ph	H
1022	5-F	CF ₃	C≡C-(3-NH ₂) Ph	H
1023	5-F	CF ₃	C≡C-(4-NH ₂) Ph	H
1024	5-F	CF ₃	C≡C-(2-NMe ₂) Ph	H
1025	5-F	CF ₃	C≡C-(3-NMe ₂) Ph	H
1026	5-F	CF ₃	C≡C-(4-NMe ₂) Ph	H
1027	5-F	CF ₃	C≡C-2-Pyridyl	H
1028	5-F	CF ₃	C≡C-2-Pyridyl	H
1029	5-F	CF ₃	C≡C-3-Pyridyl	H
1030	5-F	CF ₃	C≡C-4-Pyridyl	H
1031	5-F	CF ₃	C≡C-2-furanyl	H
1032	5-F	CF ₃	C≡C-3-furanyl	H
1033	5-F	CF ₃	C≡C-2-thienyl	H
1034	5-F	CF ₃	C≡C-3-thienyl	H
1035	5-F	CF ₃	C≡C-2-oxazolyl	H
1036	5-F	CF ₃	C≡C-2-thiazolyl	H
1037	5-F	CF ₃	C≡C-4-isoxazolyl	H
1038	5-F	CF ₃	C≡C-2-imidazolyl	H
1039	5-F	CF ₃	C=C-cycPr	H
1040	5-F	CF ₃	C=CCH ₂ CH ₂ OH	H
1041	5-F	CF ₃	C=C-CH(OH)Me	H
1042	5-F	CF ₃	C=C-Ph	H
1043	5-F	CF ₃	C=C-(2-Cl) Ph	H
1044	5-F	CF ₃	C=C-(3-Cl) Ph	H
1045	5-F	CF ₃	C=C-(4-Cl) Ph	H
1046	5-F	CF ₃	C=C-(2-F) Ph	H
1047	5-F	CF ₃	C=C-(3-F) Ph	H
1048	5-F	CF ₃	C=C-(4-F) Ph	H

1049	5-F	CF ₃	C=C-(2-OH) Ph	H
1050	5-F	CF ₃	C=C-(3-OH) Ph	H
1051	5-F	CF ₃	C=C-(4-OH) Ph	H
1052	5-F	CF ₃	C=C-(2-OMe) Ph	H
1053	5-F	CF ₃	C=C-(3-OMe) Ph	H
1054	5-F	CF ₃	C=C-(4-OMe) Ph	H
1055	5-F	CF ₃	C=C-(2-CN) Ph	H
1056	5-F	CF ₃	C=C-(3-CN) Ph	H
1057	5-F	CF ₃	C=C-(4-CN) Ph	H
1058	5-F	CF ₃	C=C-(2-NO ₂) Ph	H
1059	5-F	CF ₃	C=C-(3-NO ₂) Ph	H
1060	5-F	CF ₃	C=C-(4-NO ₂) Ph	H
1061	5-F	CF ₃	C=C-(2-NH ₂) Ph	H
1062	5-F	CF ₃	C=C-(3-NH ₂) Ph	H
1063	5-F	CF ₃	C=C-(4-NH ₂) Ph	H
1064	5-F	CF ₃	C=C-(2-NMe ₂) Ph	H
1065	5-F	CF ₃	C=C-(3-NMe ₂) Ph	H
1066	5-F	CF ₃	C=C-(4-NMe ₂) Ph	H
1067	5-F	CF ₃	C=C-2-Pyridyl	H
1068	5-F	CF ₃	C=C-3-Pyridyl	H
1069	5-F	CF ₃	C=C-4-Pyridyl	H
1070	5-F	CF ₃	C=C-2-furanyl	H
1071	5-F	CF ₃	C=C-3-furanyl	H
1072	5-F	CF ₃	C=C-2-thienyl	H
1073	5-F	CF ₃	C=C-3-thienyl	H
1074	5-F	CF ₃	C=C-2-oxazolyl	H
1075	5-F	CF ₃	C=C-2-thiazolyl	H
1076	5-F	CF ₃	C=C-4-isoxazolyl	H
1077	5-F	CF ₃	C=C-2-imidazolyl	H
1078	5-F	CF ₃	CH ₂ CH ₂ -cycPr	H
1079	5-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ OH	H
1080	5-F	CF ₃	CH ₂ CH ₂ -CH(OH)Me	H
1081	5-F	CF ₃	CH ₂ CH ₂ Ph	H
1082	5-F	CF ₃		

1083	5-F	CF ₃	CH ₂ CH ₂ -(2-Cl) Ph	H
1084	5-F	CF ₃	CH ₂ CH ₂ -(3-Cl) Ph	H
1085	5-F	CF ₃	CH ₂ CH ₂ -(4-Cl) Ph	H
1086	5-F	CF ₃	CH ₂ CH ₂ -(2-F) Ph	H
1087	5-F	CF ₃	CH ₂ CH ₂ -(3-F) Ph	H
1088	5-F	CF ₃	CH ₂ CH ₂ -(4-F) Ph	H
1089	5-F	CF ₃	CH ₂ CH ₂ -(2-OH) Ph	H
1090	5-F	CF ₃	CH ₂ CH ₂ -(3-OH) Ph	H
1091	5-F	CF ₃	CH ₂ CH ₂ -(4-OH) Ph	H
1092	5-F	CF ₃	CH ₂ CH ₂ -(2-OMe) Ph	H
1093	5-F	CF ₃	CH ₂ CH ₂ -(3-OMe) Ph	H
1094	5-F	CF ₃	CH ₂ CH ₂ -(4-OMe) Ph	H
1095	5-F	CF ₃	CH ₂ CH ₂ -(2-CN) Ph	H
1096	5-F	CF ₃	CH ₂ CH ₂ -(3-CN) Ph	H
1097	5-F	CF ₃	CH ₂ CH ₂ -(4-CN) Ph	H
1098	5-F	CF ₃	CH ₂ CH ₂ -(2-NO ₂) Ph	H
1099	5-F	CF ₃	CH ₂ CH ₂ -(3-NO ₂) Ph	H
1100	5-F	CF ₃	CH ₂ CH ₂ -(4-NO ₂) Ph	H
1101	5-F	CF ₃	CH ₂ CH ₂ -(2-NH ₂) Ph	H
1102	5-F	CF ₃	CH ₂ CH ₂ -(3-NH ₂) Ph	H
1103	5-F	CF ₃	CH ₂ CH ₂ -(4-NH ₂) Ph	H
1104	5-F	CF ₃	CH ₂ CH ₂ -(2-NMe ₂) Ph	H
1105	5-F	CF ₃	CH ₂ CH ₂ -(3-NMe ₂) Ph	H
1106	5-F	CF ₃	CH ₂ CH ₂ -(4-NMe ₂) Ph	H
1107	5-F	CF ₃	CH ₂ CH ₂ -2-Pyridyl	H
1108	5-F	CF ₃	CH ₂ CH ₂ -3-Pyridyl	H
1109	5-F	CF ₃	CH ₂ CH ₂ -4-Pyridyl	H
1110	5-F	CF ₃	CH ₂ CH ₂ -2-furanyl	H
1111	5-F	CF ₃	CH ₂ CH ₂ -3-furanyl	H
1112	5-F	CF ₃	CH ₂ CH ₂ -2-thienyl	H
1113	5-F	CF ₃	CH ₂ CH ₂ -3-thienyl	H
1114	5-F	CF ₃	CH ₂ CH ₂ -2-oxazolyl	H
1115	5-F	CF ₃	CH ₂ CH ₂ -2-thiazolyl	H
1116	5-F	CF ₃	CH ₂ CH ₂ -4-isoxazolyl	H

1117	5-F	CF ₃	CH ₂ CH ₂ -2-imidazolyl	H
1118	5-F	CF ₃	C≡C-cycPr	CH ₃
1119	5-F	CF ₃	C≡C-Ph	CH ₃
1120	5-F	CF ₃	C≡C-2-Pyridyl	CH ₃
1121	5-F	CF ₃	C≡C-3-Pyridyl	CH ₃
1122	5-F	CF ₃	C≡C-4-Pyridyl	CH ₃
1123	5-F	CF ₃	C≡C-2-furanyl	CH ₃
1124	5-F	CF ₃	C≡C-3-furanyl	CH ₃
1125	5-F	CF ₃	C≡C-2-thienyl	CH ₃
1126	5-F	CF ₃	C≡C-3-thienyl	CH ₃
1127	5-F	CF ₃	C=C-cycPr	CH ₃
1128	5-F	CF ₃	C=C-Ph	CH ₃
1129	5-F	CF ₃	C=C-2-Pyridyl	CH ₃
1130	5-F	CF ₃	C=C-3-Pyridyl	CH ₃
1131	5-F	CF ₃	C=C-4-Pyridyl	CH ₃
1132	5-F	CF ₃	C=C-2-furanyl	CH ₃
1133	5-F	CF ₃	C=C-3-furanyl	CH ₃
1134	5-F	CF ₃	C=C-2-thienyl	CH ₃
1135	5-F	CF ₃	C=C-3-thienyl	CH ₃
1136	5-F	CF ₃	CH ₂ CH ₂ -cycPr	CH ₃
1137	5-F	CF ₃	CH ₂ CH ₂ -Ph	CH ₃
1138	5-F	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₃
1139	5-F	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₃
1140	5-F	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₃
1141	5-F	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₃
1142	5-F	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₃
1143	5-F	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₃
1144	5-F	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₃
1145	5-F	CF ₃	C≡C-cycPr	CH ₂ CH ₃
1146	5-F	CF ₃	C≡C-Ph	CH ₂ CH ₃
1147	5-F	CF ₃	C≡C-2-Pyridyl	CH ₂ CH ₃
1148	5-F	CF ₃	C≡C-3-Pyridyl	CH ₂ CH ₃
1149	5-F	CF ₃	C≡C-4-Pyridyl	CH ₂ CH ₃
1150	5-F	CF ₃	C≡C-2-furanyl	CH ₂ CH ₃

1151	5-F	CF ₃	C≡C-3-furanyl	CH ₂ CH ₃
1152	5-F	CF ₃	C≡C-2-thienyl	CH ₂ CH ₃
1153	5-F	CF ₃	C≡C-3-thienyl	CH ₂ CH ₃
1154	5-F	CF ₃	C=C-cycPr	CH ₂ CH ₃
1155	5-F	CF ₃	C=C-Ph	CH ₂ CH ₃
1156	5-F	CF ₃	C=C-2-Pyridyl	CH ₂ CH ₃
1157	5-F	CF ₃	C=C-3-Pyridyl	CH ₂ CH ₃
1158	5-F	CF ₃	C=C-4-Pyridyl	CH ₂ CH ₃
1159	5-F	CF ₃	C=C-2-furanyl	CH ₂ CH ₃
1160	5-F	CF ₃	C=C-3-furanyl	CH ₂ CH ₃
1161	5-F	CF ₃	C=C-2-thienyl	CH ₂ CH ₃
1162	5-F	CF ₃	C=C-3-thienyl	CH ₂ CH ₃
1163	5-F	CF ₃	CH ₂ CH ₂ -cycPr	CH ₂ CH ₃
1164	5-F	CF ₃	CH ₂ CH ₂ -Ph	CH ₂ CH ₃
1165	5-F	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₂ CH ₃
1166	5-F	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₂ CH ₃
1167	5-F	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₂ CH ₃
1168	5-F	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₂ CH ₃
1169	5-F	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₂ CH ₃
1170	5-F	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₂ CH ₃
1171	5-F	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₂ CH ₃
1172	5-Cl, 6-F	CF ₃	C≡C-cycPr	H
1173	5-Cl, 6-F	CF ₃	C≡C-Ph	H
1174	5-Cl, 6-F	CF ₃	C≡C-2-Pyridyl	H
1175	5-Cl, 6-F	CF ₃	C≡C-3-Pyridyl	H
1176	5-Cl, 6-F	CF ₃	C≡C-4-Pyridyl	H
1177	5-Cl, 6-F	CF ₃	C≡C-2-furanyl	H
1178	5-Cl, 6-F	CF ₃	C≡C-3-furanyl	H
1179	5-Cl, 6-F	CF ₃	C≡C-2-thienyl	H
1180	5-Cl, 6-F	CF ₃	C≡C-3-thienyl	H
1181	5-Cl, 6-F	CF ₃	C=C-cycPr	H
1182	5-Cl, 6-F	CF ₃	C=C-Ph	H
1183	5-Cl, 6-F	CF ₃	C=C-2-Pyridyl	H
1184	5-Cl, 6-F	CF ₃	C=C-3-Pyridyl	H

1185	5-Cl,6-F	CF ₃	C=C-4-Pyridyl	H
1186	5-Cl,6-F	CF ₃	C=C-2-furanyl	H
1187	5-Cl,6-F	CF ₃	C=C-3-furanyl	H
1188	5-Cl,6-F	CF ₃	C=C-2-thienyl	H
1189	5-Cl,6-F	CF ₃	C=C-3-thienyl	H
1190	5-Cl,6-F	CF ₃	CH ₂ CH ₂ -cycPr	H
1191	5-Cl,6-F	CF ₃	CH ₂ CH ₂ -Ph	H
1192	5-Cl,6-F	CF ₃	CH ₂ CH ₂ -2-Pyridyl	H
1193	5-Cl,6-F	CF ₃	CH ₂ CH ₂ -3-Pyridyl	H
1194	5-Cl,6-F	CF ₃	CH ₂ CH ₂ -4-Pyridyl	H
1195	5-Cl,6-F	CF ₃	CH ₂ CH ₂ -2-furanyl	H
1196	5-Cl,6-F	CF ₃	CH ₂ CH ₂ -3-furanyl	H
1197	5-Cl,6-F	CF ₃	CH ₂ CH ₂ -2-thienyl	H
1198	5-Cl,6-F	CF ₃	CH ₂ CH ₂ -3-thienyl	H
1199	5-Cl,6-F	CF ₃	C≡C-cycPr	CH ₃
1200	5-Cl,6-F	CF ₃	C≡C-Ph	CH ₃
1201	5-Cl,6-F	CF ₃	C≡C-2-Pyridyl	CH ₃
1202	5-Cl,6-F	CF ₃	C≡C-3-Pyridyl	CH ₃
1203	5-Cl,6-F	CF ₃	C≡C-4-Pyridyl	CH ₃
1204	5-Cl,6-F	CF ₃	C≡C-2-furanyl	CH ₃
1205	5-Cl,6-F	CF ₃	C≡C-3-furanyl	CH ₃
1206	5-Cl,6-F	CF ₃	C≡C-2-thienyl	CH ₃
1207	5-Cl,6-F	CF ₃	C≡C-3-thienyl	CH ₃
1208	5-Cl,6-F	CF ₃	C=C-cycPr	CH ₃
1209	5-Cl,6-F	CF ₃	C=C-Ph	CH ₃
1210	5-Cl,6-F	CF ₃	C=C-2-Pyridyl	CH ₃
1211	5-Cl,6-F	CF ₃	C=C-3-Pyridyl	CH ₃
1212	5-Cl,6-F	CF ₃	C=C-4-Pyridyl	CH ₃
1213	5-Cl,6-F	CF ₃	C=C-2-furanyl	CH ₃
1214	5-Cl,6-F	CF ₃	C=C-3-furanyl	CH ₃
1215	5-Cl,6-F	CF ₃	C=C-2-thienyl	CH ₃
1216	5-Cl,6-F	CF ₃	C=C-3-thienyl	CH ₃
1217	5-Cl,6-F	CF ₃	CH ₂ CH ₂ -cycPr	CH ₃
1218	5-Cl,6-F	CF ₃	CH ₂ CH ₂ -Ph	CH ₃

1219	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₃
1220	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₃
1221	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₃
1222	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₃
1223	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₃
1224	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₃
1225	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₃
1226	5-F, 6-Cl	CF ₃	C≡C-cycPr	H
1227	5-F, 6-Cl	CF ₃	C≡C-Ph	H
1228	5-F, 6-Cl	CF ₃	C≡C-2-Pyridyl	H
1229	5-F, 6-Cl	CF ₃	C≡C-3-Pyridyl	H
1230	5-F, 6-Cl	CF ₃	C≡C-4-Pyridyl	H
1231	5-F, 6-Cl	CF ₃	C≡C-2-furanyl	H
1232	5-F, 6-Cl	CF ₃	C≡C-3-furanyl	H
1233	5-F, 6-Cl	CF ₃	C≡C-2-thienyl	H
1234	5-F, 6-Cl	CF ₃	C≡C-3-thienyl	H
1235	5-F, 6-Cl	CF ₃	C=C-cycPr	H
1236	5-F, 6-Cl	CF ₃	C=C-Ph	H
1237	5-F, 6-Cl	CF ₃	C=C-2-Pyridyl	H
1238	5-F, 6-Cl	CF ₃	C=C-3-Pyridyl	H
1239	5-F, 6-Cl	CF ₃	C=C-4-Pyridyl	H
1240	5-F, 6-Cl	CF ₃	C=C-2-furanyl	H
1241	5-F, 6-Cl	CF ₃	C=C-3-furanyl	H
1242	5-F, 6-Cl	CF ₃	C=C-2-thienyl	H
1243	5-F, 6-Cl	CF ₃	C=C-3-thienyl	H
1244	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -cycPr	H
1245	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -Ph	H
1246	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -2-Pyridyl	H
1247	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -3-Pyridyl	H
1248	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -4-Pyridyl	H
1249	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -2-furanyl	H
1250	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -3-furanyl	H
1251	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -2-thienyl	H
1252	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -3-thienyl	H

1253	5-F, 6-Cl	CF ₃	C≡C-cycPr	CH ₃
1254	5-F, 6-Cl	CF ₃	C≡C-Ph	CH ₃
1255	5-F, 6-Cl	CF ₃	C≡C-2-Pyridyl	CH ₃
1256	5-F, 6-Cl	CF ₃	C≡C-3-Pyridyl	CH ₃
1257	5-F, 6-Cl	CF ₃	C≡C-4-Pyridyl	CH ₃
1258	5-F, 6-Cl	CF ₃	C≡C-2-furanyl	CH ₃
1259	5-F, 6-Cl	CF ₃	C≡C-3-furanyl	CH ₃
1260	5-F, 6-Cl	CF ₃	C≡C-2-thienyl	CH ₃
1261	5-F, 6-Cl	CF ₃	C≡C-3-thienyl	CH ₃
1262	5-F, 6-Cl	CF ₃	C=C-cycPr	CH ₃
1263	5-F, 6-Cl	CF ₃	C=C-Ph	CH ₃
1264	5-F, 6-Cl	CF ₃	C=C-2-Pyridyl	CH ₃
1265	5-F, 6-Cl	CF ₃	C=C-3-Pyridyl	CH ₃
1266	5-F, 6-Cl	CF ₃	C=C-4-Pyridyl	CH ₃
1267	5-F, 6-Cl	CF ₃	C=C-2-furanyl	CH ₃
1268	5-F, 6-Cl	CF ₃	C=C-3-furanyl	CH ₃
1269	5-F, 6-Cl	CF ₃	C=C-2-thienyl	CH ₃
1270	5-F, 6-Cl	CF ₃	C=C-3-thienyl	CH ₃
1271	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -cycPr	CH ₃
1272	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -Ph	CH ₃
1273	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₃
1274	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₃
1275	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₃
1276	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₃
1277	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₃
1278	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₃
1279	5-F, 6-Cl	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₃
1280	6-Cl, 8-F	CF ₃	C≡C-cycPr	H
1281	6-Cl, 8-F	CF ₃	C≡C-Ph	H
1282	6-Cl, 8-F	CF ₃	C≡C-2-Pyridyl	H
1283	6-Cl, 8-F	CF ₃	C≡C-3-Pyridyl	H
1284	6-Cl, 8-F	CF ₃	C≡C-4-Pyridyl	H
1285	6-Cl, 8-F	CF ₃	C≡C-2-furanyl	H
1286	6-Cl, 8-F	CF ₃	C≡C-3-furanyl	H

1287	6-Cl,8-F	CF ₃	C≡C-2-thienyl	H
1288	6-Cl,8-F	CF ₃	C≡C-3-thienyl	H
1289	6-Cl,8-F	CF ₃	C=C-cycPr	H
1290	6-Cl,8-F	CF ₃	C=C-Ph	H
1291	6-Cl,8-F	CF ₃	C=C-2-Pyridyl	H
1292	6-Cl,8-F	CF ₃	C=C-3-Pyridyl	H
1293	6-Cl,8-F	CF ₃	C=C-4-Pyridyl	H
1294	6-Cl,8-F	CF ₃	C=C-2-furanyl	H
1295	6-Cl,8-F	CF ₃	C=C-3-furanyl	H
1296	6-Cl,8-F	CF ₃	C=C-2-thienyl	H
1297	6-Cl,8-F	CF ₃	C=C-3-thienyl	H
1298	6-Cl,8-F	CF ₃	CH ₂ CH ₂ -cycPr	H
1299	6-Cl,8-F	CF ₃	CH ₂ CH ₂ -Ph	H
1300	6-Cl,8-F	CF ₃	CH ₂ CH ₂ -2-Pyridyl	H
1301	6-Cl,8-F	CF ₃	CH ₂ CH ₂ -3-Pyridyl	H
1302	6-Cl,8-F	CF ₃	CH ₂ CH ₂ -4-Pyridyl	H
1303	6-Cl,8-F	CF ₃	CH ₂ CH ₂ -2-furanyl	H
1304	6-Cl,8-F	CF ₃	CH ₂ CH ₂ -3-furanyl	H
1305	6-Cl,8-F	CF ₃	CH ₂ CH ₂ -2-thienyl	H
1306	6-Cl,8-F	CF ₃	CH ₂ CH ₂ -3-thienyl	H
1307	6-Cl,8-F	CF ₃	C≡C-cycPr	CH ₃
1308	6-Cl,8-F	CF ₃	C≡C-Ph	CH ₃
1309	6-Cl,8-F	CF ₃	C≡C-2-Pyridyl	CH ₃
1310	6-Cl,8-F	CF ₃	C≡C-3-Pyridyl	CH ₃
1311	6-Cl,8-F	CF ₃	C≡C-4-Pyridyl	CH ₃
1312	6-Cl,8-F	CF ₃	C≡C-2-furanyl	CH ₃
1313	6-Cl,8-F	CF ₃	C≡C-3-furanyl	CH ₃
1314	6-Cl,8-F	CF ₃	C≡C-2-thienyl	CH ₃
1315	6-Cl,8-F	CF ₃	C≡C-3-thienyl	CH ₃
1316	6-Cl,8-F	CF ₃	C=C-cycPr	CH ₃
1317	6-Cl,8-F	CF ₃	C=C-Ph	CH ₃
1318	6-Cl,8-F	CF ₃	C=C-2-Pyridyl	CH ₃
1319	6-Cl,8-F	CF ₃	C=C-3-Pyridyl	CH ₃
1320	6-Cl,8-F	CF ₃	C=C-4-Pyridyl	CH ₃

1321	6-Cl, 8-F	CF ₃	C=C-2-furanyl	CH ₃
1322	6-Cl, 8-F	CF ₃	C=C-3-furanyl	CH ₃
1323	6-Cl, 8-F	CF ₃	C=C-2-thienyl	CH ₃
1324	6-Cl, 8-F	CF ₃	C=C-3-thienyl	CH ₃
1325	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ -cycPr	CH ₃
1326	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ -Ph	CH ₃
1327	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₃
1328	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₃
1329	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₃
1330	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₃
1331	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₃
1332	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₃
1333	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₃
1334	6-CH ₃	CF ₃	C≡C-cycPr	H
1335	6-CH ₃	CF ₃	C≡C-Ph	H
1336	6-CH ₃	CF ₃	C≡C-2-Pyridyl	H
1337	6-CH ₃	CF ₃	C≡C-3-Pyridyl	H
1338	6-CH ₃	CF ₃	C≡C-4-Pyridyl	H
1339	6-CH ₃	CF ₃	C≡C-2-furanyl	H
1340	6-CH ₃	CF ₃	C≡C-3-furanyl	H
1341	6-CH ₃	CF ₃	C≡C-2-thienyl	H
1342	6-CH ₃	CF ₃	C≡C-3-thienyl	H
1343	6-CH ₃	CF ₃	C=C-cycPr	H
1344	6-CH ₃	CF ₃	C=C-Ph	H
1345	6-CH ₃	CF ₃	C=C-2-Pyridyl	H
1346	6-CH ₃	CF ₃	C=C-3-Pyridyl	H
1347	6-CH ₃	CF ₃	C=C-4-Pyridyl	H
1348	6-CH ₃	CF ₃	C=C-2-furanyl	H
1349	6-CH ₃	CF ₃	C=C-3-furanyl	H
1350	6-CH ₃	CF ₃	C=C-2-thienyl	H
1351	6-CH ₃	CF ₃	C=C-3-thienyl	H
1352	6-CH ₃	CF ₃	CH ₂ CH ₂ -cycPr	H
1353	6-CH ₃	CF ₃	CH ₂ CH ₂ -Ph	H
1354	6-CH ₃	CF ₃	CH ₂ CH ₂ -2-Pyridyl	H

1355	6-CH ₃	CF ₃	CH ₂ CH ₂ -3-Pyridyl	H
1356	6-CH ₃	CF ₃	CH ₂ CH ₂ -4-Pyridyl	H
1357	6-CH ₃	CF ₃	CH ₂ CH ₂ -2-furanyl	H
1358	6-CH ₃	CF ₃	CH ₂ CH ₂ -3-furanyl	H
1359	6-CH ₃	CF ₃	CH ₂ CH ₂ -2-thienyl	H
1360	6-CH ₃	CF ₃	CH ₂ CH ₂ -3-thienyl	H
1361	6-CH ₃	CF ₃	C≡C-cycPr	CH ₃
1362	6-CH ₃	CF ₃	C≡C-Ph	CH ₃
1363	6-CH ₃	CF ₃	C≡C-2-Pyridyl	CH ₃
1364	6-CH ₃	CF ₃	C≡C-3-Pyridyl	CH ₃
1365	6-CH ₃	CF ₃	C≡C-4-Pyridyl	CH ₃
1366	6-CH ₃	CF ₃	C≡C-2-furanyl	CH ₃
1367	6-CH ₃	CF ₃	C≡C-3-furanyl	CH ₃
1368	6-CH ₃	CF ₃	C≡C-2-thienyl	CH ₃
1369	6-CH ₃	CF ₃	C≡C-3-thienyl	CH ₃
1370	6-CH ₃	CF ₃	C=C-cycPr	CH ₃
1371	6-CH ₃	CF ₃	C=C-Ph	CH ₃
1372	6-CH ₃	CF ₃	C=C-2-Pyridyl	CH ₃
1373	6-CH ₃	CF ₃	C=C-3-Pyridyl	CH ₃
1374	6-CH ₃	CF ₃	C=C-4-Pyridyl	CH ₃
1375	6-CH ₃	CF ₃	C=C-2-furanyl	CH ₃
1376	6-CH ₃	CF ₃	C=C-3-furanyl	CH ₃
1377	6-CH ₃	CF ₃	C=C-2-thienyl	CH ₃
1378	6-CH ₃	CF ₃	C=C-3-thienyl	CH ₃
1379	6-CH ₃	CF ₃	CH ₂ CH ₂ -cycPr	CH ₃
1380	6-CH ₃	CF ₃	CH ₂ CH ₂ -Ph	CH ₃
1381	6-CH ₃	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₃
1382	6-CH ₃	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₃
1383	6-CH ₃	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₃
1384	6-CH ₃	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₃
1385	6-CH ₃	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₃
1386	6-CH ₃	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₃
1387	6-CH ₃	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₃
1388	6-COCH ₃	CF ₃	C≡C-cycPr	H

1389	6-COCH ₃	CF ₃	C≡C-Ph	H
1390	6-COCH ₃	CF ₃	C≡C-2-Pyridyl	H
1391	6-COCH ₃	CF ₃	C≡C-3-Pyridyl	H
1392	6-COCH ₃	CF ₃	C≡C-4-Pyridyl	H
1393	6-COCH ₃	CF ₃	C≡C-2-furanyl	H
1394	6-COCH ₃	CF ₃	C≡C-3-furanyl	H
1395	6-COCH ₃	CF ₃	C≡C-2-thienyl	H
1396	6-COCH ₃	CF ₃	C≡C-3-thienyl	H
1397	6-NH ₂	CF ₃	C≡C-cycPr	H
1398	6-NH ₂	CF ₃	C≡C-Ph	H
1399	6-NH ₂	CF ₃	C≡C-2-Pyridyl	H
1400	6-NH ₂	CF ₃	C≡C-3-Pyridyl	H
1401	6-NH ₂	CF ₃	C≡C-4-Pyridyl	H
1402	6-NH ₂	CF ₃	C≡C-2-furanyl	H
1403	6-NH ₂	CF ₃	C≡C-3-furanyl	H
1404	6-NH ₂	CF ₃	C≡C-2-thienyl	H
1405	6-NH ₂	CF ₃	C≡C-3-thienyl	H
1406	6-NMe ₂	CF ₃	C≡C-cycPr	H
1407	6-NMe ₂	CF ₃	C≡C-Ph	H
1408	6-NMe ₂	CF ₃	C≡C-2-Pyridyl	H
1409	6-NMe ₂	CF ₃	C≡C-3-Pyridyl	H
1410	6-NMe ₂	CF ₃	C≡C-4-Pyridyl	H
1411	6-NMe ₂	CF ₃	C≡C-2-furanyl	H
1412	6-NMe ₂	CF ₃	C≡C-3-furanyl	H
1413	6-NMe ₂	CF ₃	C≡C-2-thienyl	H
1414	6-NMe ₂	CF ₃	C≡C-3-thienyl	H
1415	7-Cl	CF ₃	C≡C-cycPr	H
1416	7-Cl	CF ₃	C≡C-Ph	H
1417	7-Cl	CF ₃	C≡C-2-Pyridyl	H
1418	7-Cl	CF ₃	C≡C-3-Pyridyl	H
1419	7-Cl	CF ₃	C≡C-4-Pyridyl	H
1420	7-Cl	CF ₃	C≡C-2-furanyl	H
1421	7-Cl	CF ₃	C≡C-3-furanyl	H
1422	7-Cl	CF ₃	C≡C-2-thienyl	H

1423	7-Cl	CF ₃	C≡C-3-thienyl	H
1424	5,6-OCH ₂ O-	CF ₃	C≡C-cycPr	H
1425	5,6-OCH ₂ O-	CF ₃	C≡CCH ₂ CH ₂ OH	H
1426	5,6-OCH ₂ O-	CF ₃	C≡C-CH(OH)Me	H
1427	5,6-OCH ₂ O-	CF ₃	C≡C-Ph	H
1428	5,6-OCH ₂ O-	CF ₃	C≡C-(2-Cl)Ph	H
1429	5,6-OCH ₂ O-	CF ₃	C≡C-(3-Cl)Ph	H
1430	5,6-OCH ₂ O-	CF ₃	C≡C-(4-Cl)Ph	H
1431	5,6-OCH ₂ O-	CF ₃	C≡C-(2-F)Ph	H
1432	5,6-OCH ₂ O-	CF ₃	C≡C-(3-F)Ph	H
1433	5,6-OCH ₂ O-	CF ₃	C≡C-(4-F)Ph	H
1434	5,6-OCH ₂ O-	CF ₃	C≡C-(2-OH)Ph	H
1435	5,6-OCH ₂ O-	CF ₃	C≡C-(3-OH)Ph	H
1436	5,6-OCH ₂ O-	CF ₃	C≡C-(4-OH)Ph	H
1437	5,6-OCH ₂ O-	CF ₃	C≡C-(2-OMe)Ph	H
1438	5,6-OCH ₂ O-	CF ₃	C≡C-(3-OMe)Ph	H
1439	5,6-OCH ₂ O-	CF ₃	C≡C-(4-OMe)Ph	H
1440	5,6-OCH ₂ O-	CF ₃	C≡C-(2-CN)Ph	H
1441	5,6-OCH ₂ O-	CF ₃	C≡C-(3-CN)Ph	H
1442	5,6-OCH ₂ O-	CF ₃	C≡C-(4-CN)Ph	H
1443	5,6-OCH ₂ O-	CF ₃	C≡C-(2-NO ₂)Ph	H
1444	5,6-OCH ₂ O-	CF ₃	C≡C-(3-NO ₂)Ph	H
1445	5,6-OCH ₂ O-	CF ₃	C≡C-(4-NO ₂)Ph	H
1446	5,6-OCH ₂ O-	CF ₃	C≡C-(2-NH ₂)Ph	H
1447	5,6-OCH ₂ O-	CF ₃	C≡C-(3-NH ₂)Ph	H
1448	5,6-OCH ₂ O-	CF ₃	C≡C-(4-NH ₂)Ph	H
1449	5,6-OCH ₂ O-	CF ₃	C≡C-(2-NMe ₂)Ph	H
1450	5,6-OCH ₂ O-	CF ₃	C≡C-(3-NMe ₂)Ph	H
1451	5,6-OCH ₂ O-	CF ₃	C≡C-(4-NMe ₂)Ph	H
1452	5,6-OCH ₂ O-	CF ₃	C≡C-2-Pyridyl	H
1453	5,6-OCH ₂ O-	CF ₃	C≡C-2-Pyridyl	H
1454	5,6-OCH ₂ O-	CF ₃	C≡C-3-Pyridyl	H
1455	5,6-OCH ₂ O-	CF ₃	C≡C-4-Pyridyl	H
1456	5,6-OCH ₂ O-	CF ₃	C≡C-2-furanyl	H

1457	5,6-OCH ₂ O-	CF ₃	C≡C-3-furanyl	H
1458	5,6-OCH ₂ O-	CF ₃	C≡C-2-thienyl	H
1459	5,6-OCH ₂ O-	CF ₃	C≡C-3-thienyl	H
1460	5,6-OCH ₂ O-	CF ₃	C≡C-2-oxazolyl	H
1461	5,6-OCH ₂ O-	CF ₃	C≡C-2-thiazolyl	H
1462	5,6-OCH ₂ O-	CF ₃	C≡C-4-isoxazolyl	H
1463	5,6-OCH ₂ O-	CF ₃	C≡C-2-imidazolyl	H
1464	6-COCH ₃	CF ₃	C=C-cycPr	H
1465	6-COCH ₃	CF ₃	C=C-Ph	H
1466	6-COCH ₃	CF ₃	C=C-2-Pyridyl	H
1467	6-COCH ₃	CF ₃	C=C-3-Pyridyl	H
1468	6-COCH ₃	CF ₃	C=C-4-Pyridyl	H
1469	6-COCH ₃	CF ₃	C=C-2-furanyl	H
1470	6-COCH ₃	CF ₃	C=C-3-furanyl	H
1471	6-COCH ₃	CF ₃	C=C-2-thienyl	H
1472	6-COCH ₃	CF ₃	C=C-3-thienyl	H
1473	6-NH ₂	CF ₃	C=C-cycPr	H
1474	6-NH ₂	CF ₃	C=C-Ph	H
1475	6-NH ₂	CF ₃	C=C-2-Pyridyl	H
1476	6-NH ₂	CF ₃	C=C-3-Pyridyl	H
1477	6-NH ₂	CF ₃	C=C-4-Pyridyl	H
1478	6-NH ₂	CF ₃	C=C-2-furanyl	H
1479	6-NH ₂	CF ₃	C=C-3-furanyl	H
1480	6-NH ₂	CF ₃	C=C-2-thienyl	H
1481	6-NH ₂	CF ₃	C=C-3-thienyl	H
1482	6-NMe ₂	CF ₃	C=C-cycPr	H
1483	6-NMe ₂	CF ₃	C=C-Ph	H
1484	6-NMe ₂	CF ₃	C=C-2-Pyridyl	H
1485	6-NMe ₂	CF ₃	C=C-3-Pyridyl	H
1486	6-NMe ₂	CF ₃	C=C-4-Pyridyl	H
1487	6-NMe ₂	CF ₃	C=C-2-furanyl	H
1488	6-NMe ₂	CF ₃	C=C-3-furanyl	H
1489	6-NMe ₂	CF ₃	C=C-2-thienyl	H
1490	6-NMe ₂	CF ₃	C=C-3-thienyl	H

1491	7-Cl	CF ₃	C=C-cycPr	H
1492	7-Cl	CF ₃	C=C-Ph	H
1493	7-Cl	CF ₃	C=C-2-Pyridyl	H
1494	7-Cl	CF ₃	C=C-3-Pyridyl	H
1495	7-Cl	CF ₃	C=C-4-Pyridyl	H
1496	7-Cl	CF ₃	C=C-2-furanyl	H
1497	7-Cl	CF ₃	C=C-3-furanyl	H
1498	7-Cl	CF ₃	C=C-2-thienyl	H
1499	7-Cl	CF ₃	C=C-3-thienyl	H
1500	5,6-OCH ₂ O-	CF ₃	C=C-cycPr	H
1501	5,6-OCH ₂ O-	CF ₃	C=CCH ₂ CH ₂ OH	H
1502	5,6-OCH ₂ O-	CF ₃	C=C-CH(OH)Me	H
1503	5,6-OCH ₂ O-	CF ₃	C=C-Ph	H
1504	5,6-OCH ₂ O-	CF ₃	C=C-(2-Cl)Ph	H
1505	5,6-OCH ₂ O-	CF ₃	C=C-(3-Cl)Ph	H
1506	5,6-OCH ₂ O-	CF ₃	C=C-(4-Cl)Ph	H
1507	5,6-OCH ₂ O-	CF ₃	C=C-(2-F)Ph	H
1508	5,6-OCH ₂ O-	CF ₃	C=C-(3-F)Ph	H
1509	5,6-OCH ₂ O-	CF ₃	C=C-(4-F)Ph	H
1510	5,6-OCH ₂ O-	CF ₃	C=C-(2-OH)Ph	H
1511	5,6-OCH ₂ O-	CF ₃	C=C-(3-OH)Ph	H
1512	5,6-OCH ₂ O-	CF ₃	C=C-(4-OH)Ph	H
1513	5,6-OCH ₂ O-	CF ₃	C=C-(2-OMe)Ph	H
1514	5,6-OCH ₂ O-	CF ₃	C=C-(3-OMe)Ph	H
1515	5,6-OCH ₂ O-	CF ₃	C=C-(4-OMe)Ph	H
1516	5,6-OCH ₂ O-	CF ₃	C=C-(2-CN)Ph	H
1517	5,6-OCH ₂ O-	CF ₃	C=C-(3-CN)Ph	H
1518	5,6-OCH ₂ O-	CF ₃	C=C-(4-CN)Ph	H
1519	5,6-OCH ₂ O-	CF ₃	C=C-(2-NO ₂)Ph	H
1520	5,6-OCH ₂ O-	CF ₃	C=C-(3-NO ₂)Ph	H
1521	5,6-OCH ₂ O-	CF ₃	C=C-(4-NO ₂)Ph	H
1522	5,6-OCH ₂ O-	CF ₃	C=C-(2-NH ₂)Ph	H
1523	5,6-OCH ₂ O-	CF ₃	C=C-(3-NH ₂)Ph	H
1524	5,6-OCH ₂ O-	CF ₃	C=C-(4-NH ₂)Ph	H

1525	5,6-OCH ₂ O-	CF ₃	C=C-(2-NMe ₂) Ph	H
1526	5,6-OCH ₂ O-	CF ₃	C=C-(3-NMe ₂) Ph	H
1527	5,6-OCH ₂ O-	CF ₃	C=C-(4-NMe ₂) Ph	H
1528	5,6-OCH ₂ O-	CF ₃	C=C-2-Pyridyl	H
1529	5,6-OCH ₂ O-	CF ₃	C=C-2-Pyridyl	H
1530	5,6-OCH ₂ O-	CF ₃	C=C-3-Pyridyl	H
1531	5,6-OCH ₂ O-	CF ₃	C=C-4-Pyridyl	H
1532	5,6-OCH ₂ O-	CF ₃	C=C-2-furanyl	H
1533	5,6-OCH ₂ O-	CF ₃	C=C-3-furanyl	H
1534	5,6-OCH ₂ O-	CF ₃	C=C-2-thienyl	H
1535	5,6-OCH ₂ O-	CF ₃	C=C-3-thienyl	H
1536	5,6-OCH ₂ O-	CF ₃	C=C-2-oxazolyl	H
1537	5,6-OCH ₂ O-	CF ₃	C=C-2-thiazolyl	H
1538	5,6-OCH ₂ O-	CF ₃	C=C-4-isoxazolyl	H
1539	5,6-OCH ₂ O-	CF ₃	C=C-2-imidazolyl	H
1540	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -cycPr	H
1541	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ OH	H
1542	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -CH(OH) Me	H
1543	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ Ph	H
1544	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(2-Cl) Ph	H
1545	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(3-Cl) Ph	H
1546	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(4-Cl) Ph	H
1547	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(2-F) Ph	H
1548	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(3-F) Ph	H
1549	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(4-F) Ph	H
1550	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(2-OH) Ph	H
1551	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(3-OH) Ph	H
1552	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(4-OH) Ph	H
1553	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(2-OMe) Ph	H
1554	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(3-OMe) Ph	H
1555	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(4-OMe) Ph	H
1556	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(2-CN) Ph	H
1557	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(3-CN) Ph	H
1558	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(4-CN) Ph	H


1559	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(2-NO ₂) Ph	H
1560	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(3-NO ₂) Ph	H
1561	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(4-NO ₂) Ph	H
1562	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(2-NH ₂) Ph	H
1563	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(3-NH ₂) Ph	H
1564	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(4-NH ₂) Ph	H
1565	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(2-NMe ₂) Ph	H
1566	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(3-NMe ₂) Ph	H
1567	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -(4-NMe ₂) Ph	H
1568	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -2-Pyridyl	H
1569	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -3-Pyridyl	H
1570	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -4-Pyridyl	H
1571	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -2-furanyl	H
1572	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -3-furanyl	H
1573	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -2-thienyl	H
1574	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -3-thienyl	H
1575	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -2-oxazolyl	H
1576	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -2-thiazolyl	H
1577	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -4-isoxazolyl	H
1578	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -2-imidazolyl	H
1579	5,6-OCH ₂ O-	CF ₃	C≡C-cycPr	CH ₃
1580	5,6-OCH ₂ O-	CF ₃	C≡C-Ph	CH ₃
1581	5,6-OCH ₂ O-	CF ₃	C≡C-2-Pyridyl	CH ₃
1582	5,6-OCH ₂ O-	CF ₃	C≡C-3-Pyridyl	CH ₃
1583	5,6-OCH ₂ O-	CF ₃	C≡C-4-Pyridyl	CH ₃
1584	5,6-OCH ₂ O-	CF ₃	C≡C-2-furanyl	CH ₃
1585	5,6-OCH ₂ O-	CF ₃	C≡C-3-furanyl	CH ₃
1586	5,6-OCH ₂ O-	CF ₃	C≡C-2-thienyl	CH ₃
1587	5,6-OCH ₂ O-	CF ₃	C≡C-3-thienyl	CH ₃
1588	5,6-OCH ₂ O-	CF ₃	C=C-cycPr	CH ₃
1589	5,6-OCH ₂ O-	CF ₃	C=C-Ph	CH ₃
1590	5,6-OCH ₂ O-	CF ₃	C=C-2-Pyridyl	CH ₃
1591	5,6-OCH ₂ O-	CF ₃	C=C-3-Pyridyl	CH ₃
1592	5,6-OCH ₂ O-	CF ₃	C=C-4-Pyridyl	CH ₃

1593	5,6-OCH ₂ O-	CF ₃	C=C-2-furanyl	CH ₃
1594	5,6-OCH ₂ O-	CF ₃	C=C-3-furanyl	CH ₃
1595	5,6-OCH ₂ O-	CF ₃	C=C-2-thienyl	CH ₃
1596	5,6-OCH ₂ O-	CF ₃	C=C-3-thienyl	CH ₃
1597	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -cycPr	CH ₃
1598	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -Ph	CH ₃
1599	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₃
1600	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₃
1601	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₃
1602	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₃
1603	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₃
1604	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₃
1605	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₃
1606	5,6-OCH ₂ O-	CF ₃	C≡C-cycPr	CH ₂ CH ₃
1607	5,6-OCH ₂ O-	CF ₃	C≡C-Ph	CH ₂ CH ₃
1608	5,6-OCH ₂ O-	CF ₃	C≡C-2-Pyridyl	CH ₂ CH ₃
1609	5,6-OCH ₂ O-	CF ₃	C≡C-3-Pyridyl	CH ₂ CH ₃
1610	5,6-OCH ₂ O-	CF ₃	C≡C-4-Pyridyl	CH ₂ CH ₃
1611	5,6-OCH ₂ O-	CF ₃	C≡C-2-furanyl	CH ₂ CH ₃
1612	5,6-OCH ₂ O-	CF ₃	C≡C-3-furanyl	CH ₂ CH ₃
1613	5,6-OCH ₂ O-	CF ₃	C≡C-2-thienyl	CH ₂ CH ₃
1614	5,6-OCH ₂ O-	CF ₃	C≡C-3-thienyl	CH ₂ CH ₃
1615	5,6-OCH ₂ O-	CF ₃	C=C-cycPr	CH ₂ CH ₃
1616	5,6-OCH ₂ O-	CF ₃	C=C-Ph	CH ₂ CH ₃
1617	5,6-OCH ₂ O-	CF ₃	C=C-2-Pyridyl	CH ₂ CH ₃
1618	5,6-OCH ₂ O-	CF ₃	C=C-3-Pyridyl	CH ₂ CH ₃
1619	5,6-OCH ₂ O-	CF ₃	C=C-4-Pyridyl	CH ₂ CH ₃
1620	5,6-OCH ₂ O-	CF ₃	C=C-2-furanyl	CH ₂ CH ₃
1621	5,6-OCH ₂ O-	CF ₃	C=C-3-furanyl	CH ₂ CH ₃
1622	5,6-OCH ₂ O-	CF ₃	C=C-2-thienyl	CH ₂ CH ₃
1623	5,6-OCH ₂ O-	CF ₃	C=C-3-thienyl	CH ₂ CH ₃
1624	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -cycPr	CH ₂ CH ₃
1625	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -Ph	CH ₂ CH ₃
1626	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -2-Pyridyl	CH ₂ CH ₃

1627	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -3-Pyridyl	CH ₂ CH ₃
1628	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -4-Pyridyl	CH ₂ CH ₃
1629	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -2-furanyl	CH ₂ CH ₃
1630	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -3-furanyl	CH ₂ CH ₃
1631	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -2-thienyl	CH ₂ CH ₃
1632	5,6-OCH ₂ O-	CF ₃	CH ₂ CH ₂ -3-thienyl	CH ₂ CH ₃

*Unless otherwise indicated, stereochemistry is (+/-).

Table 3*



Ex. #	R ³	R ¹	R ²	R ⁸
1	6-Cl	CF ₃	C≡C-Pr	H
2	6-Cl	CF ₃	C≡C-Bu	H
3	6-Cl	CF ₃	C≡C-iBu	H
4	6-Cl	CF ₃	C≡C-tBu	H
5	6-Cl	CF ₃	C≡C-Me	H
6	6-Cl	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	H
7	6-Cl	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	H
8	6-Cl	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	H
9	6-Cl	CF ₃	CH ₂ CH ₂ CH ₃	H
10	6-Cl	CF ₃	CH ₂ CH ₂ -tBu	H
11	6-Cl	CF ₃	CH ₂ C≡C-CH ₃	H
12	6-Cl	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	H
13	6-Cl	CF ₃	C≡C-iPr	CH ₃
14	6-Cl	CF ₃	C≡C-Pr	CH ₃
15	6-Cl	CF ₃	C≡C-Bu	CH ₃
16	6-Cl	CF ₃	C≡C-iBu	CH ₃
17	6-Cl	CF ₃	C≡C-tBu	CH ₃
18	6-Cl	CF ₃	C≡C-Et	CH ₃
19	6-Cl	CF ₃	C≡C-Me	CH ₃
20	6-Cl	CF ₃	CH ₂ C≡C-CH ₃	CH ₃
21	6-Cl	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₃
22	6-Cl	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₃
23	6-Cl	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₃
24	6-Cl	CF ₃	CH ₂ CH ₂ CH ₃	CH ₃
25	6-Cl	CF ₃	CH ₂ CH ₂ -tBu	CH ₃
26	6-Cl	CF ₃	C≡C-iPr	CH ₂ CH ₃
27	6-Cl	CF ₃	C≡C-Pr	CH ₂ CH ₃
28	6-Cl	CF ₃	C≡C-Bu	CH ₂ CH ₃

29	6-Cl	CF ₃	C≡C-iBu	CH ₂ CH ₃
30	6-Cl	CF ₃	C≡C-tBu	CH ₂ CH ₃
31	6-Cl	CF ₃	C≡C-Et	CH ₂ CH ₃
32	6-Cl	CF ₃	C≡C-Me	CH ₂ CH ₃
33	6-Cl	CF ₃	CH ₂ C≡C-CH ₃	CH ₂ CH ₃
34	6-Cl	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₂ CH ₃
35	6-Cl	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₂ CH ₃
36	6-Cl	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
37	6-Cl	CF ₃	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
38	6-Cl	CF ₃	CH ₂ CH ₂ -tBu	CH ₂ CH ₃
39	6-MeO	CF ₃	C≡C-Pr	H
40	6-MeO	CF ₃	C≡C-Bu	H
41	6-MeO	CF ₃	C≡C-iBu	H
42	6-MeO	CF ₃	C≡C-tBu	H
43	6-MeO	CF ₃	C≡C-Et	H
44	6-MeO	CF ₃	C≡C-Me	H
45	6-MeO	CF ₃	CH ₂ C≡C-CH ₃	H
46	6-MeO	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	H
47	6-MeO	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	H
48	6-MeO	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	H
49	6-MeO	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	H
50	6-MeO	CF ₃	CH ₂ CH ₂ CH ₃	H
51	6-MeO	CF ₃	CH ₂ CH ₂ -tBu	H
52	6-MeO	CF ₃	CH ₂ C≡C-CH ₃	H
53	6-MeO	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	H
54	6-MeO	CF ₃	C≡C-iPr	CH ₃
55	6-MeO	CF ₃	C≡C-Pr	CH ₃
56	6-MeO	CF ₃	C≡C-Bu	CH ₃
57	6-MeO	CF ₃	C≡C-iBu	CH ₃
58	6-MeO	CF ₃	C≡C-tBu	CH ₃
59	6-MeO	CF ₃	C≡C-Et	CH ₃
60	6-MeO	CF ₃	C≡C-Me	CH ₃
61	6-MeO	CF ₃	CH ₂ C≡C-CH ₃	CH ₃
62	6-MeO	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₃

63	6-MeO	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₃
64	6-MeO	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₃
65	6-MeO	CF ₃	CH ₂ CH ₂ CH ₃	CH ₃
66	6-MeO	CF ₃	CH ₂ CH ₂ -tBu	CH ₃
67	6-MeO	CF ₃	C≡C-iPr	CH ₂ CH ₃
68	6-MeO	CF ₃	C≡C-Pr	CH ₂ CH ₃
69	6-MeO	CF ₃	C≡C-Bu	CH ₂ CH ₃
70	6-MeO	CF ₃	C≡C-iBu	CH ₂ CH ₃
71	6-MeO	CF ₃	C≡C-tBu	CH ₂ CH ₃
72	6-MeO	CF ₃	C≡C-Et	CH ₂ CH ₃
73	6-MeO	CF ₃	C≡C-Me	CH ₂ CH ₃
74	6-MeO	CF ₃	CH ₂ C≡C-CH ₃	CH ₂ CH ₃
75	6-MeO	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₂ CH ₃
76	6-MeO	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₂ CH ₃
77	6-MeO	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
78	6-MeO	CF ₃	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
79	6-MeO	CF ₃	CH ₂ CH ₂ -tBu	CH ₂ CH ₃
80	5,6-diF	CF ₃	C≡C-Pr	H
81	5,6-diF	CF ₃	C≡C-Bu	H
82	5,6-diF	CF ₃	C≡C-iBu	H
83	5,6-diF	CF ₃	C≡C-tBu	H
84	5,6-diF	CF ₃	C≡C-Me	H
85	5,6-diF	CF ₃	CH ₂ C≡C-CH ₃	H
86	5,6-diF	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	H
87	5,6-diF	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	H
88	5,6-diF	CF ₃	CH ₂ CH ₂ CH ₃	H
89	5,6-diF	CF ₃	CH ₂ CH ₂ -tBu	H
90	5,6-diF	CF ₃	C≡C-iPr	CH ₃
91	5,6-diF	CF ₃	C≡C-Pr	CH ₃
92	5,6-diF	CF ₃	C≡C-Bu	CH ₃
93	5,6-diF	CF ₃	C≡C-iBu	CH ₃
94	5,6-diF	CF ₃	C≡C-tBu	CH ₃
95	5,6-diF	CF ₃	C≡C-Et	CH ₃
96	5,6-diF	CF ₃	C≡C-Me	CH ₃

97	5,6-diF	CF ₃	C≡C-Ph	CH ₃
98	5,6-diF	CF ₃	CH ₂ C≡C-CH ₃	CH ₃
99	5,6-diF	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₃
100	5,6-diF	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₃
101	5,6-diF	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₃
102	5,6-diF	CF ₃	CH ₂ CH ₂ CH ₃	CH ₃
103	5,6-diF	CF ₃	CH ₂ CH ₂ -tBu	CH ₃
104	5,6-diF	CF ₃	C≡C-iPr	CH ₂ CH ₃
105	5,6-diF	CF ₃	C≡C-Pr	CH ₂ CH ₃
106	5,6-diF	CF ₃	C≡C-Bu	CH ₂ CH ₃
107	5,6-diF	CF ₃	C≡C-iBu	CH ₂ CH ₃
108	5,6-diF	CF ₃	C≡C-tBu	CH ₂ CH ₃
109	5,6-diF	CF ₃	C≡C-Et	CH ₂ CH ₃
110	5,6-diF	CF ₃	C≡C-Me	CH ₂ CH ₃
111	5,6-diF	CF ₃	CH ₂ C≡C-CH ₃	CH ₂ CH ₃
112	5,6-diF	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₂ CH ₃
113	5,6-diF	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₂ CH ₃
114	5,6-diF	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
115	5,6-diF	CF ₃	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
116	5,6-diF	CF ₃	CH ₂ CH ₂ -tBu	CH ₂ CH ₃
117	6-F	CF ₃	C≡C-Pr	H
118	6-F	CF ₃	C≡C-Bu	H
119	6-F	CF ₃	C≡C-iBu	H
120	6-F	CF ₃	C≡C-tBu	H
121	6-F	CF ₃	C≡C-Me	H
122	6-F	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	H
123	6-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	H
124	6-F	CF ₃	CH ₂ CH ₂ CH ₃	H
125	6-F	CF ₃	CH ₂ CH ₂ -tBu	H
126	6-F	CF ₃	C≡C-iPr	CH ₃
127	6-F	CF ₃	C≡C-Pr	CH ₃
128	6-F	CF ₃	C≡C-Bu	CH ₃
129	6-F	CF ₃	C≡C-iBu	CH ₃
130	6-F	CF ₃	C≡C-tBu	CH ₃

131	6-F	CF ₃	C≡C-Et	CH ₃
132	6-F	CF ₃	C≡C-Me	CH ₃
133	6-F	CF ₃	CH ₂ C≡C-CH ₃	CH ₃
134	6-F	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₃
135	6-F	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₃
136	6-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₃
137	6-F	CF ₃	CH ₂ CH ₂ CH ₃	CH ₃
138	6-F	CF ₃	CH ₂ CH ₂ -tBu	CH ₃
139	6-F	CF ₃	C≡C-iPr	CH ₂ CH ₃
140	6-F	CF ₃	C≡C-Pr	CH ₂ CH ₃
141	6-F	CF ₃	C≡C-Bu	CH ₂ CH ₃
142	6-F	CF ₃	C≡C-iBu	CH ₂ CH ₃
143	6-F	CF ₃	C≡C-tBu	CH ₂ CH ₃
144	6-F	CF ₃	C≡C-Et	CH ₂ CH ₃
145	6-F	CF ₃	C≡C-Me	CH ₂ CH ₃
146	6-F	CF ₃	CH ₂ C≡C-CH ₃	CH ₂ CH ₃
147	6-F	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₂ CH ₃
148	6-F	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₂ CH ₃
149	6-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
150	6-F	CF ₃	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
151	6-F	CF ₃	CH ₂ CH ₂ -tBu	CH ₂ CH ₃
152	5-Cl	CF ₃	C≡C-iPr	H
153	5-Cl	CF ₃	C≡C-Pr	H
154	5-Cl	CF ₃	C≡C-Bu	H
155	5-Cl	CF ₃	C≡C-iBu	H
156	5-Cl	CF ₃	C≡C-tBu	H
157	5-Cl	CF ₃	C≡C-Et	H
158	5-Cl	CF ₃	C≡C-Me	H
159	5-Cl	CF ₃	CH ₂ C≡C-CH ₃	H
160	5-Cl	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	H
161	5-Cl	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	H
162	5-Cl	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	H
163	5-Cl	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	H
164	5-Cl	CF ₃	CH ₂ CH ₂ CH ₃	H

165	5-Cl	CF ₃	CH ₂ CH ₂ -tBu	H
166	5-Cl	CF ₃	C≡C-iPr	CH ₃
167	5-Cl	CF ₃	C≡C-Pr	CH ₃
168	5-Cl	CF ₃	C≡C-Bu	CH ₃
169	5-Cl	CF ₃	C≡C-iBu	CH ₃
170	5-Cl	CF ₃	C≡C-tBu	CH ₃
171	5-Cl	CF ₃	C≡C-Et	CH ₃
172	5-Cl	CF ₃	C≡C-Me	CH ₃
173	5-Cl	CF ₃	CH ₂ C≡C-CH ₃	CH ₃
174	5-Cl	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₃
175	5-Cl	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₃
176	5-Cl	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₃
177	5-Cl	CF ₃	CH ₂ CH ₂ CH ₃	CH ₃
178	5-Cl	CF ₃	CH ₂ CH ₂ -tBu	CH ₃
179	5-Cl	CF ₃	C≡C-iPr	CH ₂ CH ₃
180	5-Cl	CF ₃	C≡C-Pr	CH ₂ CH ₃
181	5-Cl	CF ₃	C≡C-Bu	CH ₂ CH ₃
182	5-Cl	CF ₃	C≡C-iBu	CH ₂ CH ₃
183	5-Cl	CF ₃	C≡C-tBu	CH ₂ CH ₃
184	5-Cl	CF ₃	C≡C-Et	CH ₂ CH ₃
185	5-Cl	CF ₃	C≡C-Me	CH ₂ CH ₃
186	5-Cl	CF ₃	CH ₂ C≡C-CH ₃	CH ₂ CH ₃
187	5-Cl	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₂ CH ₃
188	5-Cl	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₂ CH ₃
189	5-Cl	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
190	5-Cl	CF ₃	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
191	5-Cl	CF ₃	CH ₂ CH ₂ -tBu	CH ₂ CH ₃
192	5-F	CF ₃	C≡C-iPr	H
193	5-F	CF ₃	C≡C-Pr	H
194	5-F	CF ₃	C≡C-Bu	H
195	5-F	CF ₃	C≡C-iBu	H
196	5-F	CF ₃	C≡C-tBu	H
197	5-F	CF ₃	C≡C-Et	H
198	5-F	CF ₃	C≡C-Me	H

199	5-F	CF ₃	CH ₂ C≡C-CH ₃	H
200	5-F	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	H
201	5-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	H
202	5-F	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	H
203	5-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	H
204	5-F	CF ₃	CH ₂ CH ₂ CH ₃	H
205	5-F	CF ₃	CH ₂ CH ₂ -tBu	H
206	5-F	CF ₃	C≡C-iPr	CH ₃
207	5-F	CF ₃	C≡C-Pr	CH ₃
208	5-F	CF ₃	C≡C-Bu	CH ₃
209	5-F	CF ₃	C≡C-iBu	CH ₃
210	5-F	CF ₃	C≡C-tBu	CH ₃
211	5-F	CF ₃	C≡C-Et	CH ₃
212	5-F	CF ₃	C≡C-Me	CH ₃
213	5-F	CF ₃	CH ₂ C≡C-CH ₃	CH ₃
214	5-F	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₃
215	5-F	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₃
216	5-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₃
217	5-F	CF ₃	CH ₂ CH ₂ CH ₃	CH ₃
218	5-F	CF ₃	CH ₂ CH ₂ -tBu	CH ₃
219	5-F	CF ₃	C≡C-iPr	CH ₂ CH ₃
220	5-F	CF ₃	C≡C-Pr	CH ₂ CH ₃
221	5-F	CF ₃	C≡C-Bu	CH ₂ CH ₃
222	5-F	CF ₃	C≡C-iBu	CH ₂ CH ₃
223	5-F	CF ₃	C≡C-tBu	CH ₂ CH ₃
224	5-F	CF ₃	C≡C-Et	CH ₂ CH ₃
225	5-F	CF ₃	C≡C-Me	CH ₂ CH ₃
226	5-F	CF ₃	CH ₂ C≡C-CH ₃	CH ₂ CH ₃
227	5-F	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₂ CH ₃
228	5-F	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₂ CH ₃
229	5-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
230	5-F	CF ₃	CH ₂ CH ₂ CH ₃	CH ₂ CH ₃
231	5-F	CF ₃	CH ₂ CH ₂ -tBu	CH ₂ CH ₃
232	5-Cl, 6-F	CF ₃	C≡C-iPr	H

233	5-Cl, 6-F	CF ₃	C≡C-Pr	H
234	5-Cl, 6-F	CF ₃	C≡C-Bu	H
235	5-Cl, 6-F	CF ₃	C≡C-iBu	H
236	5-Cl, 6-F	CF ₃	C≡C-tBu	H
237	5-Cl, 6-F	CF ₃	C≡C-Et	H
238	5-Cl, 6-F	CF ₃	C≡C-Me	H
239	5-Cl, 6-F	CF ₃	CH ₂ C≡C-CH ₃	H
240	5-Cl, 6-F	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	H
241	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	H
242	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	H
243	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ CH ₃	H
244	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ -tBu	H
245	5-Cl, 6-F	CF ₃	C≡C-iPr	CH ₃
246	5-Cl, 6-F	CF ₃	C≡C-Pr	CH ₃
247	5-Cl, 6-F	CF ₃	C≡C-Bu	CH ₃
248	5-Cl, 6-F	CF ₃	C≡C-iBu	CH ₃
249	5-Cl, 6-F	CF ₃	C≡C-tBu	CH ₃
250	5-Cl, 6-F	CF ₃	C≡C-Et	CH ₃
251	5-Cl, 6-F	CF ₃	C≡C-Me	CH ₃
252	5-Cl, 6-F	CF ₃	CH ₂ C≡C-CH ₃	CH ₃
253	5-Cl, 6-F	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₃
254	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₃
255	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₃
256	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ CH ₃	CH ₃
257	5-Cl, 6-F	CF ₃	CH ₂ CH ₂ -tBu	CH ₃
258	6-Cl, 8-F	CF ₃	C≡C-iPr	H
259	6-Cl, 8-F	CF ₃	C≡C-Pr	H
260	6-Cl, 8-F	CF ₃	C≡C-Bu	H
261	6-Cl, 8-F	CF ₃	C≡C-iBu	H
262	6-Cl, 8-F	CF ₃	C≡C-tBu	H
263	6-Cl, 8-F	CF ₃	C≡C-Et	H
264	6-Cl, 8-F	CF ₃	C≡C-Me	H
265	6-Cl, 8-F	CF ₃	CH ₂ C≡C-CH ₃	H
266	6-Cl, 8-F	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	H

267	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	H
268	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	H
269	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ CH ₃	H
270	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ -tBu	H
271	6-Cl, 8-F	CF ₃	C≡C-iPr	CH ₃
272	6-Cl, 8-F	CF ₃	C≡C-Pr	CH ₃
273	6-Cl, 8-F	CF ₃	C≡C-Bu	CH ₃
274	6-Cl, 8-F	CF ₃	C≡C-iBu	CH ₃
275	6-Cl, 8-F	CF ₃	C≡C-tBu	CH ₃
276	6-Cl, 8-F	CF ₃	C≡C-Et	CH ₃
277	6-Cl, 8-F	CF ₃	C≡C-Me	CH ₃
278	6-Cl, 8-F	CF ₃	CH ₂ C≡C-CH ₃	CH ₃
279	6-Cl, 8-F	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₃
280	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₃
281	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₃
282	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ CH ₃	CH ₃
283	6-Cl, 8-F	CF ₃	CH ₂ CH ₂ -tBu	CH ₃
284	6-CH ₃	CF ₃	C≡C-iPr	H
285	6-CH ₃	CF ₃	C≡C-Pr	H
286	6-CH ₃	CF ₃	C≡C-Bu	H
287	6-CH ₃	CF ₃	C≡C-iBu	H
288	6-CH ₃	CF ₃	C≡C-tBu	H
289	6-CH ₃	CF ₃	C≡C-Et	H
290	6-CH ₃	CF ₃	C≡C-Me	H
291	6-CH ₃	CF ₃	CH ₂ C≡C-CH ₃	H
292	6-CH ₃	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	H
293	6-CH ₃	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	H
294	6-CH ₃	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	H
295	6-CH ₃	CF ₃	CH ₂ CH ₂ CH ₃	H
296	6-CH ₃	CF ₃	CH ₂ CH ₂ -tBu	H
297	6-CH ₃	CF ₃	C≡C-iPr	CH ₃
298	6-CH ₃	CF ₃	C≡C-Pr	CH ₃
299	6-CH ₃	CF ₃	C≡C-Bu	CH ₃
300	6-CH ₃	CF ₃	C≡C-iBu	CH ₃

301	6-CH ₃	CF ₃	C≡C-tBu	CH ₃
302	6-CH ₃	CF ₃	C≡C-Et	CH ₃
303	6-CH ₃	CF ₃	C≡C-Me	CH ₃
304	6-CH ₃	CF ₃	CH ₂ C≡C-CH ₃	CH ₃
305	6-CH ₃	CF ₃	CH ₂ C≡C-CH ₂ CH ₃	CH ₃
306	6-CH ₃	CF ₃	CH ₂ CH ₂ CH(CH ₃) ₂	CH ₃
307	6-CH ₃	CF ₃	CH ₂ CH ₂ CH ₂ CH ₃	CH ₃
308	6-CH ₃	CF ₃	CH ₂ CH ₂ CH ₃	CH ₃
309	6-CH ₃	CF ₃	CH ₂ CH ₂ -tBu	CH ₃
310	6-COCH ₃	CF ₃	C≡C-iPr	H
311	6-COCH ₃	CF ₃	C≡C-Pr	H
312	6-COCH ₃	CF ₃	C≡C-Bu	H
313	6-COCH ₃	CF ₃	C≡C-iBu	H
314	6-COCH ₃	CF ₃	C≡C-tBu	H
315	6-COCH ₃	CF ₃	C≡C-Et	H
316	6-COCH ₃	CF ₃	C≡C-Me	H
317	6-NH ₂	CF ₃	C≡C-iPr	H
318	6-NH ₂	CF ₃	C≡C-Pr	H
319	6-NH ₂	CF ₃	C≡C-Bu	H
320	6-NH ₂	CF ₃	C≡C-iBu	H
321	6-NH ₂	CF ₃	C≡C-tBu	H
322	6-NH ₂	CF ₃	C≡C-Et	H
323	6-NH ₂	CF ₃	C≡C-Me	H
324	6-NMe ₂	CF ₃	C≡C-iPr	H
325	6-NMe ₂	CF ₃	C≡C-Pr	H
326	6-NMe ₂	CF ₃	C≡C-Bu	H
327	6-NMe ₂	CF ₃	C≡C-iBu	H
328	6-NMe ₂	CF ₃	C≡C-tBu	H
329	6-NMe ₂	CF ₃	C≡C-Et	H
330	6-NMe ₂	CF ₃	C≡C-Me	H
331	7-Cl	CF ₃	C≡C-iPr	H
332	7-Cl	CF ₃	C≡C-Pr	H
333	7-Cl	CF ₃	C≡C-Bu	H
334	7-Cl	CF ₃	C≡C-iBu	H

335	7-Cl	CF ₃	C≡C-tBu	H
336	7-Cl	CF ₃	C≡C-Et	H
337	7-Cl	CF ₃	C≡C-Me	H

*Unless otherwise indicated, stereochemistry is (+/-).

Utility

The compounds of this invention possess reverse transcriptase inhibitory activity, in particular, HIV inhibitory efficacy. The compounds of formula (I) possess HIV reverse transcriptase inhibitory activity and are therefore useful as antiviral agents for the treatment of HIV infection and associated diseases. The compounds of formula (I) possess HIV reverse transcriptase inhibitory activity and are effective as inhibitors of HIV growth. The ability of the compounds of the present invention to inhibit viral growth or infectivity is demonstrated in standard assay of viral growth or infectivity, for example, using the assay described below.

The compounds of formula (I) of the present invention are also useful for the inhibition of HIV in an ex vivo sample containing HIV or expected to be exposed to HIV. Thus, the compounds of the present invention may be used to inhibit HIV present in a body fluid sample (for example, a serum or semen sample) which contains or is suspected to contain or be exposed to HIV.

The compounds provided by this invention are also useful as standard or reference compounds for use in tests or assays for determining the ability of an agent to inhibit viral clone replication and/or HIV reverse transcriptase, for example in a pharmaceutical research program. Thus, the compounds of the present invention may be used as a control or reference compound in such assays and as a quality control standard. The compounds of the present invention may be provided in a commercial kit or container for use as such standard or reference compound.

Since the compounds of the present invention exhibit specificity for HIV reverse transcriptase, the compounds of the present invention may also be useful as diagnostic reagents in diagnostic assays for the detection of HIV reverse transcriptase. Thus, inhibition of the reverse transcriptase activity in an assay (such as the assays described herein) by a compound of the present invention

would be indicative of the presence of HIV reverse transcriptase and HIV virus.

As used herein "µg" denotes microgram, "mg" denotes milligram, "g" denotes gram, "µL" denotes microliter, "mL" denotes milliliter, "L" denotes liter, "nM" denotes nanomolar, "µM" denotes micromolar, "mM" denotes millimolar, "M" denotes molar and "nm" denotes nanometer. "Sigma" stands for the Sigma-Aldrich Corp. of St. Louis, MO.

10 HIV RNA Assay

DNA Plasmids and in vitro RNA transcripts:

Plasmid pDAB 72 containing both gag and pol sequences of BH10 (bp 113-1816) cloned into PTZ 19R was prepared according to Erickson-Viitanen et al. *AIDS Research and Human Retroviruses* **1989**, 5, 577. The plasmid was linearized with Bam HI prior to the generation of in vitro RNA transcripts using the Riboprobe Gemini system II kit (Promega) with T7 RNA polymerase. Synthesized RNA was purified by treatment with RNase free DNase (Promega), phenol-chloroform extraction, and ethanol precipitation. RNA transcripts were dissolved in water, and stored at -70°C. The concentration of RNA was determined from the A₂₆₀.

25 Probes:

Biotinylated capture probes were purified by HPLC after synthesis on an Applied Biosystems (Foster City, CA) DNA synthesizer by addition of biotin to the 5' terminal end of the oligonucleotide, using the biotin-phosphoramidite reagent of Cocuzza, *Tet. Lett.* **1989**, 30, 6287. The gag biotinylated capture probe (5-biotin-CTAGCTCCCTGCTTGCCCATACTA 3') was complementary to nucleotides 889-912 of HXB2 and the pol biotinylated capture probe (5'-biotin -CCCTATCATTTTTGGTTTCCAT 3') was complementary to nucleotides 2374-2395 of HXB2. Alkaline phosphatase conjugated oligonucleotides used as reporter probes were prepared by Syngene (San Diego, CA.). The pol reporter probe (5' CTGTCTTACTTTGATAAAACCTC 3') was complementary to nucleotides 2403-2425 of HXB2. The gag

reporter probe (5' CCCAGTATTTGTCTACAGCCTTCT 3') was complementary to nucleotides 950-973 of HXB2. All nucleotide positions are those of the GenBank Genetic Sequence Data Bank as accessed through the Genetics Computer Group Sequence Analysis Software Package (Devereau *Nucleic Acids Research* 1984, 12, 387). The reporter probes were prepared as 0.5 μ M stocks in 2 x SSC (0.3 M NaCl, 0.03 M sodium citrate), 0.05 M Tris pH 8.8, 1 mg/mL BSA. The biotinylated capture probes were prepared as 100 μ M stocks in water.

10

Streptavidin coated plates:

Streptavidin coated plates were obtained from Du Pont Biotechnology Systems (Boston, MA).

15 Cells and virus stocks:

MT-2 and MT-4 cells were maintained in RPMI 1640 supplemented with 5% fetal calf serum (FCS) for MT-2 cells or 10% FCS for MT-4 cells, 2 mM L-glutamine and 50 μ g/mL gentamycin, all from Gibco. HIV-1 RF was propagated in MT-4 cells in the same medium. Virus stocks were prepared approximately 10 days after acute infection of MT-4 cells and stored as aliquots at -70°C. Infectious titers of HIV-1(RF) stocks were 1-3 x 10⁷ PFU (plaque forming units)/mL as measured by plaque assay on MT-2 cells (see below). Each aliquot of virus stock used for infection was thawed only once.

For evaluation of antiviral efficacy, cells to be infected were subcultured one day prior to infection. On the day of infection, cells were resuspended at 5 x 10⁵ cells/mL in RPMI 1640, 5% FCS for bulk infections or at 2 x 10⁶/mL in Dulbecco's modified Eagles medium with 5% FCS for infection in microtiter plates. Virus was added and culture continued for 3 days at 37°C.

35 HIV RNA assay:

Cell lysates or purified RNA in 3 M or 5 M GED were mixed with 5 M GED and capture probe to a final guanidinium isothiocyanate concentration of 3 M and a final biotin

oligonucleotide concentration of 30 nM. Hybridization was carried out in sealed U bottom 96 well tissue culture plates (Nunc or Costar) for 16-20 hours at 37°C. RNA hybridization reactions were diluted three-fold with deionized water to a final guanidinium isothiocyanate concentration of 1 M and aliquots (150 µL) were transferred to streptavidin coated microtiter plates wells. Binding of capture probe and capture probe-RNA hybrid to the immobilized streptavidin was allowed to proceed for 2 hours at room temperature, after which the plates were washed 6 times with DuPont ELISA plate wash buffer (phosphate buffered saline(PBS), 0.05% Tween 20.) A second hybridization of reporter probe to the immobilized complex of capture probe and hybridized target RNA was carried out in the washed streptavidin coated well by addition of 120 µl of a hybridization cocktail containing 4 X SSC, 0.66% Triton X 100, 6.66% deionized formamide, 1 mg/mL BSA and 5 nM reporter probe. After hybridization for one hour at 37°C, the plate was again washed 6 times. Immobilized alkaline phosphatase activity was detected by addition of 100 µL of 0.2 mM 4-methylumbelliferyl phosphate (MUBP, JBL Scientific) in buffer δ (2.5 M diethanolamine pH 8.9 (JBL Scientific), 10 mM MgCl₂, 5 mM zinc acetate dihydrate and 5 mM N-hydroxyethyl-ethylene-diamine-triacetic acid). The plates were incubated at 37°C. Fluorescence at 450 nM was measured using a microplate fluorometer (Dynateck) exciting at 365 nM.

Microplate based compound evaluation in HIV-1 infected MT-2 cells:

Compounds to be evaluated were dissolved in DMSO and diluted in culture medium to twice the highest concentration to be tested and a maximum DMSO concentration of 2%. Further three-fold serial dilutions of the compound in culture medium were performed directly in U bottom microtiter plates (Nunc). After compound dilution, MT-2 cells (50 µL) were added to a final concentration of 5×10^5 per mL (1×10^5 per well). Cells were incubated with compounds for 30 minutes at 37°C in a CO₂ incubator. For evaluation of antiviral potency, an

appropriate dilution of HIV-1 (RF) virus stock (50 μ L) was added to culture wells containing cells and dilutions of the test compounds. The final volume in each well was 200 μ L. Eight wells per plate were left uninfected with 50 μ L of medium added in place of virus, while eight wells were infected in the absence of any antiviral compound. For evaluation of compound toxicity, parallel plates were cultured without virus infection.

After 3 days of culture at 37°C in a humidified chamber inside a CO₂ incubator, all but 25 μ L of medium/well was removed from the HIV infected plates. Thirty seven μ L of 5 M GED containing biotinylated capture probe was added to the settled cells and remaining medium in each well to a final concentration of 3 M GED and 30 nM capture probe.

Hybridization of the capture probe to HIV RNA in the cell lysate was carried out in the same microplate well used for virus culture by sealing the plate with a plate sealer (Costar), and incubating for 16-20 hrs in a 37°C incubator. Distilled water was then added to each well to dilute the hybridization reaction three-fold and 150 μ L of this diluted mixture was transferred to a streptavidin coated microtiter plate. HIV RNA was quantitated as described above. A standard curve, prepared by adding known amounts of pDAB 72 in vitro RNA transcript to wells containing lysed uninfected cells, was run on each microtiter plate in order to determine the amount of viral RNA made during the infection.

In order to standardize the virus inoculum used in the evaluation of compounds for antiviral activity, dilutions of virus were selected which resulted in an IC₉₀ value

(concentration of compound required to reduce the HIV RNA level by 90%) for dideoxycytidine (ddC) of 0.2 μ g/mL. IC₉₀ values of other antiviral compounds, both more and less potent than ddC, were reproducible using several stocks of HIV-1 (RF) when this procedure was followed. This concentration of virus corresponded to $\sim 3 \times 10^5$ PFU (measured by plaque assay on MT-2 cells) per assay well and typically produced approximately 75% of the maximum viral RNA level achievable at any virus inoculum. For the HIV RNA assay,

IC₉₀ values were determined from the percent reduction of net signal (signal from infected cell samples minus signal from uninfected cell samples) in the RNA assay relative to the net signal from infected, untreated cells on the same culture plate (average of eight wells). Valid performance of individual infection and RNA assay tests was judged according to three criteria. It was required that the virus infection should result in an RNA assay signal equal to or greater than the signal generated from 2 ng of pDAB 72 *in vitro* RNA transcript. The IC₉₀ for ddC, determined in each assay run, should be between 0.1 and 0.3 µg/mL. Finally, the plateau level of viral RNA produced by an effective reverse transcriptase inhibitor should be less than 10% of the level achieved in an uninhibited infection. A compound was considered active if its IC₉₀ was found to be less than 20µM.

For antiviral potency tests, all manipulations in microtiter plates, following the initial addition of 2X concentrated compound solution to a single row of wells, were performed using a Perkin Elmer/Cetus ProPette.

Protein Binding and Mutant Resistance

In order to characterize NNRTI analogs for their clinical efficacy potential the effect of plasma proteins on antiviral potency and measurements of antiviral potency against wild type and mutant variants of HIV which carry amino acid changes in the known binding site for NNRTIs were examined. The rationale for this testing strategy is two fold:

1. Many drugs are extensively bound to plasma proteins. Although the binding affinity for most drugs for the major components of human plasma, namely, human serum albumin (HSA) or alpha-1-acid glycoprotein (AAG), is low, these major components are present in high concentration in the blood. Only free or unbound drug is available to cross the infected cell membrane for interaction with the target site (i.e., HIV-1 reverse transcriptase, HIV-1 RT). Therefore, the effect of added HSA+AAG on the antiviral potency in tissue culture more closely reflects the potency of a given compound

in the clinical setting. The concentration of compound required for 90% inhibition of virus replication as measured in a sensitive viral RNA-based detection method is designated the IC90. The fold increase in apparent IC90 for test compounds in the presence or added levels of HSA and AAG that reflect *in vivo* concentrations (45 mg/ml HSA, 1 mg/ml AAG) was then calculated. The lower the fold increase, the more compound will be available to interact with the target site.

2. The combination of the high rate of virus replication in the infected individual and the poor fidelity of the viral RT results in the production of a quasi-species or mixtures of HIV species in the infected individual. These species will include a majority wild type species, but also mutant variants of HIV and the proportion of a given mutant will reflect its relative fitness and replication rate. Because mutant variants including mutants with changes in the amino acid sequence of the viral RT likely pre-exist in the infected individual's quasi-species, the overall potency observed in the clinical setting will reflect the ability of a drug to inhibit not only wild type HIV-1, but mutant variants as well. We thus have constructed, in a known genetic background, mutant variants of HIV-1 which carry amino acid substitutions at positions thought to be involved in NNRTI binding, and measured the ability of test compounds to inhibit replication of these mutant viruses. The concentration of compound required for 90% inhibition of virus replication as measured in a sensitive viral RNA-based detection method is designated the IC90. It is desirable to have a compound which has high activity against a variety of mutants.

Dosage and Formulation

The antiviral compounds of this invention can be administered as treatment for viral infections by any means that produces contact of the active agent with the agent's site of action, i.e., the viral reverse transcriptase, in the body of a mammal. They can be administered by any conventional means available for use in conjunction with

pharmaceuticals, either as individual therapeutic agents or in a combination of therapeutic agents. They can be administered alone, but preferably are administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

The dosage administered will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration; the age, health and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; and the effect desired. A daily dosage of active ingredient can be expected to be about 0.001 to about 1000 milligrams per kilogram of body weight, with the preferred dose being about 0.1 to about 30 mg/kg.

Dosage forms of compositions suitable for administration contain from about 1 mg to about 100 mg of active ingredient per unit. In these pharmaceutical compositions the active ingredient will ordinarily be present in an amount of about 0.5-95% by weight based on the total weight of the composition. The active ingredient can be administered orally in solid dosage forms, such as capsules, tablets and powders, or in liquid dosage forms, such as elixirs, syrups and suspensions. It can also be administered parenterally, in sterile liquid dosage forms.

Gelatin capsules contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract. Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts, and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propyl-paraben and chlorobutanol. Suitable pharmaceutical carriers are described in *Remington's Pharmaceutical Sciences, supra*, a standard reference text in this field.

Useful pharmaceutical dosage-forms for administration of the compounds of this invention can be illustrated as follows:

20 Capsules

A large number of unit capsules can be prepared by filling standard two-piece hard gelatin capsules each with 100 mg of powdered active ingredient, 150 mg of lactose, 50 mg of cellulose, and 6 mg magnesium stearic.

25

Soft Gelatin Capsules

A mixture of active ingredient in a digestible oil such as soybean oil, cottonseed oil or olive oil can be prepared and injected by means of a positive displacement pump into gelatin to form soft gelatin capsules containing 100 mg of the active ingredient. The capsules should then be washed and dried.

Tablets

35 A large number of tablets can be prepared by conventional procedures so that the dosage unit is 100 mg of active ingredient, 0.2 mg of colloidal silicon dioxide, 5 milligrams of magnesium stearate, 275 mg of microcrystalline

cellulose, 11 mg of starch and 98.8 mg of lactose. Appropriate coatings may be applied to increase palatability or delay absorption.

5 Suspension

 An aqueous suspension can be prepared for oral administration so that each 5 mL contain 25 mg of finely divided active ingredient, 200 mg of sodium carboxymethyl cellulose, 5 mg of sodium benzoate, 1.0 g of sorbitol
10 solution, U.S.P., and 0.025 mg of vanillin.

Injectable

 A parenteral composition suitable for administration by injection can be prepared by stirring 1.5% by weight of
15 active ingredient in 10% by volume propylene glycol and water. The solution is sterilized by commonly used techniques.

Combination of components (a) and (b)

20 Each therapeutic agent component of this invention can independently be in any dosage form, such as those described above, and can also be administered in various ways, as described above. In the following description component (b)
25 is to be understood to represent one or more agents as described previously. Thus, if components (a) and (b) are to be treated the same or independently, each agent of component (b) may also be treated the same or independently.

 Components (a) and (b) of the present invention may be formulated together, in a single dosage unit (that is,
30 combined together in one capsule, tablet, powder, or liquid, etc.) as a combination product. When component (a) and (b) are not formulated together in a single dosage unit, the component (a) may be administered at the same time as component (b) or in any order; for example component (a) of
35 this invention may be administered first, followed by administration of component (b), or they may be administered in the reverse order. If component (b) contains more than one agent, e.g., one RT inhibitor and one protease inhibitor,

these agents may be administered together or in any order. When not administered at the same time, preferably the administration of component (a) and (b) occurs less than about one hour apart. Preferably, the route of
5 administration of component (a) and (b) is oral. The terms oral agent, oral inhibitor, oral compound, or the like, as used herein, denote compounds which may be orally administered. Although it is preferable that component (a) and component (b) both be administered by the same route
10 (that is, for example, both orally) or dosage form, if desired, they may each be administered by different routes (that is, for example, one component of the combination product may be administered orally, and another component may be administered intravenously) or dosage forms.

15 As is appreciated by a medical practitioner skilled in the art, the dosage of the combination therapy of the invention may vary depending upon various factors such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration, the age, health and
20 weight of the recipient, the nature and extent of the symptoms, the kind of concurrent treatment, the frequency of treatment, and the effect desired, as described above.

The proper dosage of components (a) and (b) of the present invention will be readily ascertainable by a medical
25 practitioner skilled in the art, based upon the present disclosure. By way of general guidance, typically a daily dosage may be about 100 milligrams to about 1.5 grams of each component. If component (b) represents more than one compound, then typically a daily dosage may be about 100
30 milligrams to about 1.5 grams of each agent of component (b). By way of general guidance, when the compounds of component (a) and component (b) are administered in combination, the dosage amount of each component may be reduced by about 70-80% relative to the usual dosage of the component when it is
35 administered alone as a single agent for the treatment of HIV infection, in view of the synergistic effect of the combination.

The combination products of this invention may be formulated such that, although the active ingredients are combined in a single dosage unit, the physical contact between the active ingredients is minimized. In order to minimize contact, for example, where the product is orally administered, one active ingredient may be enteric coated. By enteric coating one of the active ingredients, it is possible not only to minimize the contact between the combined active ingredients, but also, it is possible to control the release of one of these components in the gastrointestinal tract such that one of these components is not released in the stomach but rather is released in the intestines. Another embodiment of this invention where oral administration is desired provides for a combination product wherein one of the active ingredients is coated with a sustained-release material which effects a sustained-release throughout the gastrointestinal tract and also serves to minimize physical contact between the combined active ingredients. Furthermore, the sustained-released component can be additionally enteric coated such that the release of this component occurs only in the intestine. Still another approach would involve the formulation of a combination product in which the one component is coated with a sustained and/or enteric release polymer, and the other component is also coated with a polymer such as a lowviscosity grade of hydroxypropyl methylcellulose or other appropriate materials as known in the art, in order to further separate the active components. The polymer coating serves to form an additional barrier to interaction with the other component. In each formulation wherein contact is prevented between components (a) and (b) via a coating or some other material, contact may also be prevented between the individual agents of component (b).

Dosage forms of the combination products of the present invention wherein one active ingredient is enteric coated can be in the form of tablets such that the enteric coated component and the other active ingredient are blended together and then compressed into a tablet or such that the

enteric coated component is compressed into one tablet layer and the other active ingredient is compressed into an additional layer. Optionally, in order to further separate the two layers, one or more placebo layers may be present
5 such that the placebo layer is between the layers of active ingredients. In addition, dosage forms of the present invention can be in the form of capsules wherein one active ingredient is compressed into a tablet or in the form of a plurality of microtablets, particles, granules or non-perils,
10 which are then enteric coated. These enteric coated microtablets, particles, granules or non-perils are then placed into a capsule or compressed into a capsule along with a granulation of the other active ingredient.

These as well as other ways of minimizing contact
15 between the components of combination products of the present invention, whether administered in a single dosage form or administered in separate forms but at the same time or concurrently by the same manner, will be readily apparent to those skilled in the art, based on the present disclosure.


20 Pharmaceutical kits useful for the treatment of HIV infection, which comprise a therapeutically effective amount of a pharmaceutical composition comprising a compound of component (a) and one or more compounds of component (b), in one or more sterile containers, are also within the ambit of
25 the present invention. Sterilization of the container may be carried out using conventional sterilization methodology well known to those skilled in the art. Component (a) and component (b) may be in the same sterile container or in separate sterile containers. The sterile containers of
30 materials may comprise separate containers, or one or more multi-part containers, as desired. Component (a) and component (b), may be separate, or physically combined into a single dosage form or unit as described above. Such kits may further include, if desired, one or more of various
35 conventional pharmaceutical kit components, such as for example, one or more pharmaceutically acceptable carriers, additional vials for mixing the components, etc., as will be readily apparent to those skilled in the art. Instructions,

either as inserts or as labels, indicating quantities of the components to be administered, guidelines for administration, and/or guidelines for mixing the components, may also be included in the kit.

- 5 Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described herein.

WHAT IS CLAIMED AS NEW AND DESIRED TO BE SECURED BY
LETTER PATENT OF UNITED STATES IS:

1. A compound of formula (I):



I

or a stereoisomer or pharmaceutically acceptable salt thereof, wherein:

R¹ is C₁₋₃ alkyl substituted with 1-7 halogen;

R² is selected from C₁₋₅ alkyl substituted with 1-2 R⁴, C₂₋₅ alkenyl substituted with 1-2 R⁴, and C₂₋₅ alkynyl substituted with 1 R⁴;

R³, at each occurrence, is independently selected from C₁₋₄ alkyl, OH, C₁₋₄ alkoxy, F, Cl, Br, I, NR⁵R^{5a}, NO₂, CN, C(O)R⁶, NHC(O)R⁷, and NHC(O)NR⁵R^{5a};

alternatively, if two R³'s are present and are attached to adjacent carbons, then they may combine to form -OCH₂O-;

R⁴ is selected from C₃₋₅ cycloalkyl substituted with 0-2 R³, phenyl substituted with 0-5 R³, and a 5-6 membered heterocyclic system containing 1-3 heteroatoms selected from O, N, and S, substituted with 0-2 R³;

R⁵ and R^{5a} are independently selected from H and C₁₋₃ alkyl;

R⁶ is selected from H, OH, C₁₋₄ alkyl, C₁₋₄ alkoxy, and NR⁵R^{5a};

R⁷ is selected from C₁₋₃ alkyl and C₁₋₃ alkoxy;

R⁸ is selected from H, C₃₋₅ cycloalkyl, and C₁₋₃ alkyl; and,

n is selected from 0, 1, 2, 3, and 4.

5 2. A compound according to Claim 1, wherein:

R¹ is C₁₋₃ alkyl substituted with 1-7 halogen;

10 R² is selected from C₁₋₅ alkyl substituted with 1 R⁴, C₂₋₅
alkenyl substituted with 1 R⁴, and C₂₋₅ alkynyl
substituted with 1 R⁴;

15 R³, at each occurrence, is independently selected from C₁₋₄
alkyl, OH, C₁₋₄ alkoxy, F, Cl, Br, I, NR⁵R^{5a}, NO₂, CN,
C(O)R⁶, NHC(O)R⁷, and NHC(O)NR⁵R^{5a};

alternatively, if two R³'s are present and are attached to
adjacent carbons, then they may combine to form -OCH₂O-;

20 R⁴ is selected from C₃₋₅ cycloalkyl substituted with 0-2 R³,
phenyl substituted with 0-2 R³, and a 5-6 membered
heterocyclic system containing 1-3 heteroatoms selected
from O, N, and S, substituted with 0-1 R³;

25 R⁵ and R^{5a} are independently selected from H, CH₃ and C₂H₅;

R⁶ is selected from H, OH, CH₃, C₂H₅, OCH₃, OC₂H₅, and NR⁵R^{5a};

30 R⁷ is selected from CH₃, C₂H₅, OCH₃, and OC₂H₅;

R⁸ is selected from H, cyclopropyl, CH₃ and C₂H₅; and,

n is selected from 0, 1, 2, and 3.

35

3. A compound according to Claim 2, wherein:

R¹ is selected from CF₃, and C₂F₅;

R² is selected from C₁₋₃ alkyl substituted with 1 R⁴, C₂₋₃ alkenyl substituted with 1 R⁴, and C₂₋₃ alkynyl substituted with 1 R⁴;

5

R³, at each occurrence, is independently selected from C₁₋₃ alkyl, OH, C₁₋₃ alkoxy, F, Cl, Br, I, NR⁵R^{5a}, NO₂, CN, C(O)R⁶, NHC(O)R⁷, and NHC(O)NR⁵R^{5a};

10 alternatively, if two R³'s are present and are attached to adjacent carbons, then they may combine to form -OCH₂O-;

R⁴ is selected from C₃₋₅ cycloalkyl substituted with 0-2 R³, phenyl substituted with 0-2 R³, and a 5-6 membered
15 heterocyclic system containing 1-3 heteroatoms selected from O, N, and S, substituted with 0-1 R³;

R⁵ and R^{5a} are independently selected from H, CH₃ and C₂H₅;

20 R⁶ is selected from H, OH, CH₃, C₂H₅, OCH₃, OC₂H₅, and NR⁵R^{5a};

R⁷ is selected from CH₃, C₂H₅, OCH₃, and OC₂H₅;

R⁸ is selected from H, CH₃ and C₂H₅; and,

25

n is selected from 0, 1, and 2.

4. A compound according to Claim 3, wherein:

30

R¹ is CF₃;

R² is selected from C₁₋₃ alkyl substituted with 1 R⁴, C₂₋₃ alkenyl substituted with 1 R⁴, and C₂₋₃ alkynyl
35 substituted with 1 R⁴;

R^3 , at each occurrence, is independently selected from C_{1-3} alkyl, OH, C_{1-3} alkoxy, F, Cl, NR^5R^{5a} , NO_2 , CN, $C(O)R^6$, $NHC(O)R^7$, and $NHC(O)NR^5R^{5a}$;

5 alternatively, if two R^3 's are present and are attached to adjacent carbons, then they may combine to form $-OCH_2O-$;

R^4 is selected from cyclopropyl substituted with 0-1 R^3 , phenyl substituted with 0-2 R^3 , and a 5-6 membered
10 heterocyclic system containing 1-3 heteroatoms selected from O, N, and S, substituted with 0-1 R^3 , wherein the heterocyclic system is selected from 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-furanyl, 3-furanyl, 2-thienyl, 3-thienyl, 2-oxazolyl, 2-thiazolyl, 4-isoxazolyl, and 2-
15 imidazolyl;

R^5 and R^{5a} are independently selected from H, CH_3 and C_2H_5 ;

R^6 is selected from H, OH, CH_3 , C_2H_5 , OCH_3 , OC_2H_5 , and NR^5R^{5a} ;


20

R^7 is selected from CH_3 , C_2H_5 , OCH_3 , and OC_2H_5 ;

R^8 is selected from H, CH_3 and C_2H_5 ; and,

25 n is selected from 1 and 2.


5. A compound according to Claim 4, wherein the compound is of formula Ia:



30

Ia.

6. A compound according to Claim 4, wherein the
35 compound is of formula Ia:



Ib.

5 7. A compound according to Claim 1, wherein the compound is selected from:

(+/-)-6-Chloro-4-cyclopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

10

(+/-)-6-Chloro-4-(2-pyridyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

15

(+/-)-6-Chloro-4-phenylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

(+/-)-4-Cyclopropylethynyl-6-methoxy-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

20

(+/-)-6-Methoxy-4-(2-pyridyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

(+/-)-6-Methoxy-4-phenylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

25

(+/-)-4-Cyclopropylethynyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

30

(+/-)-5,6-Difluoro-4-(2-pyridyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

(+/-)-5,6-Difluoro-4-phenylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

(+/-)-4-Cyclopropylethynyl-6-fluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

5 (+/-)-6-Fluoro-4-(2-pyridyl)ethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

(+/-)-6-Fluoro-4-phenylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

10 (+/-)-6-Fluoro-4-(2'-2-pyridyl)ethyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

(+/-)-6-Fluoro-4-phenylethyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

15 (-)-6-Chloro-4-cyclopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

20 (+)-6-Chloro-4-cyclopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

(+)-4-Cyclopropylethynyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

25 (-)-4-Cyclopropylethynyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;


(+)-E-4-Cyclopropylethenyl-5,6-difluoro-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone; and,

30 (-)-6-Chloro-4-E-cyclopropylethenyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

or a pharmaceutically acceptable salt thereof.

35

8. A compound according of formula II:



II

or a stereoisomer or pharmaceutically acceptable salt thereof, wherein:

5

R^2 is $C\equiv C-R^{4a}$;

R^3 is selected from C_{1-4} alkyl, OH, C_{1-4} alkoxy, F, Cl, Br, I, NR^5R^{5a} , NO_2 , CN, $C(O)R^6$, $NHC(O)R^7$, and $NHC(O)NR^5R^{5a}$;

10

R^{4a} is selected from methyl, ethyl, n-propyl, i-propyl, i-butyl, t-butyl, and i-pentyl;

R^5 and R^{5a} are independently selected from H and C_{1-3} alkyl;

15

R^6 is selected from H, OH, C_{1-4} alkyl, C_{1-4} alkoxy, and NR^5R^{5a} ;

R^7 is selected from C_{1-3} alkyl and C_{1-3} alkoxy;

20

R^8 is selected from H, C_{3-5} cycloalkyl, and C_{1-3} alkyl; and,

n is selected from 0, 1, 2, 3, and 4.

25

9. A compound according to Claim 8, wherein:

R^2 is $C\equiv C-R^{4a}$;

R^3 is selected from C_{1-4} alkyl, OH, C_{1-4} alkoxy, F, Cl, Br, I, NR^5R^{5a} , NO_2 , CN, $C(O)R^6$, and $NHC(O)R^7$;

30

R^{4a} is selected from methyl, ethyl, n-propyl, i-propyl, i-butyl, t-butyl, and i-pentyl;

35

R^5 and R^{5a} are independently selected from H, CH_3 and C_2H_5 ;


R⁶ is selected from H, OH, CH₃, C₂H₅, OCH₃, OC₂H₅, and NR⁵R^{5a};

R⁷ is selected from CH₃, C₂H₅, OCH₃, and OC₂H₅;

R⁸ is selected from H, cyclopropyl, CH₃ and C₂H₅; and,


n is selected from 0, 1, and 2.

10. A compound according to Claim 9, wherein the compound is of formula IIa:



IIa.

11. A compound according to Claim 9, wherein the compound is of formula IIb:



IIb.

12. A compound according to Claim 8, wherein the compound is selected from:

(+/-)-6-Chloro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

(+/-)-6-Chloro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;

- (+/-)-4-Isopropylethynyl-6-methoxy-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (+/-)-5,6-Difluoro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (+/-)-5,6-Difluoro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (+/-)-5,6-Difluoro-4-isopentyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (+/-)-6-Fluoro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (+/-)-6-Fluoro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (-)-5,6-Difluoro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (+)-5,6-Difluoro-4-isopropylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- (-)-5,6-Difluoro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone; and,
- (+)-5,6-Difluoro-4-ethylethynyl-4-trifluoromethyl-3,4-dihydro-2(1H)-quinazolinone;
- or a pharmaceutically acceptable salt thereof.

13. A pharmaceutical composition, comprising: a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of one of Claims 1-12 or a pharmaceutically acceptable salt form thereof.

14. A method for treating HIV infection, comprising:
administering to a host in need of such treatment a
therapeutically effective amount of a compound of one of
5 Claims 1-12, or a pharmaceutically acceptable salt form
thereof.

15. A method of treating HIV infection which comprises
10 administering, in combination, to a host in need thereof a
therapeutically effective amount of:

(a) a compound of one of Claims 1-12 or stereoisomeric
forms, mixtures of stereoisomeric forms, or pharmaceutically
acceptable salts thereof; and,

15 (b) at least one compound selected from the group
consisting of HIV reverse transcriptase inhibitors and HIV
protease inhibitors.

20 16. A method according to Claim 15, wherein the reverse
transcriptase inhibitor is selected from AZT, 3TC, ddI, ddC,
d4T, delavirdine, TIBO derivatives, BI-RG-587, nevirapine, L-
697,661, LY 73497, Ro 18,893, loviride, trovirdine, MKC-442,
and HBY 097, and the protease inhibitor is selected from
25 saquinavir, ritonavir, indinavir, VX-478, nelfinavir, KNI-
272, CGP-61755, U-140690, and ABT-378.

30 17. A method according to Claim 16, wherein the reverse
transcriptase inhibitor is selected from AZT and 3TC and the
protease inhibitor is selected from saquinavir, nelfinavir,
ritonavir, and indinavir.

35 18. A pharmaceutical kit useful for the treatment of
HIV infection, which comprises a therapeutically effective
amount of:

(a) a compound of one of Claims 1-12 or stereoisomeric forms, mixtures of stereoisomeric forms, or pharmaceutically acceptable salts thereof; and,

(b) at least one compound selected from the group
5 consisting of HIV reverse transcriptase inhibitors and HIV protease inhibitors, in one or more sterile containers.


THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 239/80, 401/06, 405/06, 409/06, 413/06, 417/06, 403/06, A61K 31/505	A3	(11) International Publication Number: WO 98/45276 (43) International Publication Date: 15 October 1998 (15.10.98)
(21) International Application Number: PCT/US98/06733 (22) International Filing Date: 7 April 1998 (07.04.98) (30) Priority Data: 08/838,540 9 April 1997 (09.04.97) US 60/071,322 14 January 1998 (14.01.98) US (71) Applicant: DU PONT PHARMACEUTICALS COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors: CORBETT, Jeffrey, W.; 1639 Oak Avenue, Haddon Height, NJ 08035 (US). KO, Soo, Sung; 7 Aston Circle, Hockessin, DE 19707 (US). (74) Agent: VANCE, David, H.; Du Pont Pharmaceuticals Com- pany, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		(81) Designated States: AM, AU, AZ, BR, BY, CA, CN, CZ, EE, HU, IL, JP, KG, KR, KZ, LT, LV, MD, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, UA, VN, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> (88) Date of publication of the international search report: 14 January 1999 (14.01.99)

(54) Title: 4,4-DISUBSTITUTED-3,4-DIHYDRO-2(1H)-QUINAZOLINONES USEFUL AS HIV REVERSE TRANSCRIPTASE INHIBITORS

**(57) Abstract**

The present invention relates to 4,4-disubstituted-3,4-dihydro-2(1H)-quinazolines of formula (I), or stereoisomeric forms, stereoisomeric mixtures, or pharmaceutically acceptable salt forms thereof, which are useful as inhibitors of HIV reverse transcriptase, and to pharmaceutical compositions and diagnostic kits comprising the same, and methods of using the same for treating viral infection or as an assay standard or reagent.

THIS PAGE BLANK (USPTO)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/06733

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D239/80 C07D401/06 C07D405/06 C07D409/06 C07D413/06
C07D417/06 C07D403/06 A61K31/505

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 530 994 A (MERCK & CO. INC.) 10 March 1993 cited in the application see page 37, line 30 - page 45, line 54; claims 1,8-10; tables I,II & WO 93 04047 A cited in the application ---	1-6,8, 13,18
Y	DE 43 20 347 A (BOEHRINGER MANNHEIM GMBH) 22 December 1994 cited in the application see claims 1,8,10,11; example 2 --- -/--	1-6,8,13

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"&" document member of the same patent family

Date of the actual completion of the international search

14 October 1998

Date of mailing of the international search report

03. 11. 98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Hass, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/06733

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 569 083 A (MERCK & CO. INC.) 10 November 1993 see pages 4, 5, table see page 11, line 35 - page 20, line 58; claims 1,8-10 & WO 95 12583 A cited in the application ---	1-6,13, 18
Y	US 5 519 021 A (S. D. YOUNG ET AL.) 21 May 1996 cited in the application see column 3, compounds 37.2, 4, 2; table I, compounds 1 to 4, 6 to 12, 16 see claims 1,2,4,5 ---	1-6,13, 18
Y	T. J. TUCKER ET AL.: JOURNAL OF MEDICINAL CHEMISTRY, vol. 37, 1994, pages 2437-44, XP002072024 cited in the application see page 2437, abstract; page 2439, table 2; page 2438, scheme 2 ---	1,5,6
A	I. N. HOUPIS ET AL.: TETRAHEDRON LETTERS, vol. 35, no. 37, 1994, pages 6811-4, XP002072025 cited in the application see scheme III ---	1,5
A	M. A. HUFFMAN ET AL.: JOURNAL OF ORGANIC CHEMISTRY, vol. 60, 1995, pages 1590-4, XP002072026 cited in the application see page 1591, scheme 1 ---	1-6
A	WO 95 13273 A (MERCK & CO., INC.) 18 May 1995 cited in the application see claim 1; examples ---	1-6
A	GB 2 281 297 A (MERCK & CO INC) 1 March 1995 see claim 1; examples ---	1-6
A	T. PRUEKSARITANONT ET AL.: DRUG METABOLISM AND DISPOSITION, vol. 23, no. 7, 1995, pages 688-95, XP002072027 see page 689, figure 1 ---	1-6
A	M. YAMAMOTO ET AL.: CHEMICAL AND PHARMACEUTICAL BULLETIN, vol. 29, no. 8, 1981, pages 2135-56, XP002080827 see page 2139, table III -----	8

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/ 06733

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 14-17
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 14-17 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition. Rule 39.1(iv) PCT.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. Claims: 1-7, 13 (partly), 18 (partly)
2. Claims: 8-12, 13 (partly), 18 (partly)

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/06733

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 530994	A	10-03-1993	AU 2436792 A CN 1071917 A MX 9204720 A WO 9304047 A	16-03-1993 12-05-1993 01-07-1993 04-03-1993
DE 4320347	A	22-12-1994	NONE	
EP 569083	A	10-11-1993	AU 3841393 A AU 4220493 A BG 99149 A CA 2095194 A CN 1085550 A CZ 9402724 A EP 0639184 A FI 945199 A HR 930857 A HU 71401 A JP 2097872 C JP 6009578 A JP 8013805 B MX 9302681 A NO 944208 A SI 9300244 A WO 9322292 A ZA 9303179 A	11-11-1993 29-11-1993 30-06-1995 08-11-1993 20-04-1994 18-10-1995 22-02-1995 04-11-1994 30-06-1995 28-11-1995 02-10-1996 18-01-1994 14-02-1996 31-05-1994 06-01-1995 31-12-1993 11-11-1993 07-11-1994
US 5519021	A	21-05-1996	AU 1694295 A WO 9520389 A US 5663169 A US 5665720 A US 5811423 A AU 4449693 A BG 99383 A CA 2101572 A CN 1090277 A CZ 9500286 A EP 0582455 A FI 950508 A HR 931102 A HU 71219 A	15-08-1995 03-08-1995 02-09-1997 09-09-1997 22-09-1998 10-02-1994 29-09-1995 08-02-1994 03-08-1994 13-09-1995 09-02-1994 06-02-1995 31-10-1997 28-11-1995

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/06733

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5519021 A		IL 106507 A	20-11-1997
		JP 2063220 C	24-06-1996
		JP 6184124 A	05-07-1994
		JP 7080860 B	30-08-1995
		MX 9304775 A	31-05-1994
		NO 950424 A	06-04-1995
		NZ 255216 A	28-05-1996
		PL 307348 A	15-05-1995
		SI 9300419 A	30-06-1994
		SK 16195 A	11-07-1995
		WO 9403440 A	17-02-1994
		ZA 9305724 A	03-03-1994
WO 9513273 A	18-05-1995	US 5434152 A	18-07-1995
		AU 1048695 A	29-05-1995
		HR 940906 A	31-12-1996
GB 2281297 A	01-03-1995	US 5457201 A	10-10-1995

THIS PAGE BLANK (USPTO)